

This application is a Continuation-in-Part of PCT/GB00/01822, filed May 12, 2000, designating the U.S., published November 23, 2000 as WO 00/69886 and claiming priority from GB 9911298.9 filed May 14, 1999 and GB 9930530.2 filed December 23, 1999. All of the above-mentioned applications, as well as all documents cited herein and documents referenced or cited in documents cited herein, are hereby incorporated herein by reference.

10007334-10801

## Oligomeric Chaperone Proteins

The present invention relates to oligomers of chaperone or minichaperone proteins. In particular, the invention relates to polypeptides constructed by oligomerisation of  
5 minichaperone monomers to form a heptameric ring structure.

Proteins, especially catalytic proteins (enzymes) and proteins which have biological activities, are dependent on tertiary structure for most or all of their functional attributes. Tertiary structure, which is defined by the three-dimensional arrangement of the protein,  
10 is dependent on the folding of the primary polypeptide sequence in three dimensions. The tertiary structure is stabilised by the interaction between parts of the primary sequence in the folded state, such as the formation of disulphide bonds, and energetic considerations deriving from juxtaposition of particular chemical entities in the three-dimensional arrangement.

15 It is known that the tertiary structure of proteins, particularly proteins which are stored for any length of time, can degrade, leading to suboptimal activity. This can be due to a variety of factors, including aggregation and the formation of improper intra- and inter-molecular interactions. Moreover, it is also known that proteins produced by recombinant  
20 DNA technology are frequently misfolded, especially if produced in bacterial expression systems. The strongly reducing environment present in bacterial cytoplasm impedes the formation of correct disulphide bonds, thus obstructing the folding process.

Many proteins require the assistance of molecular chaperones in order to be fold *in vivo*  
25 or to be refolded *in vitro* in high yields. Molecular chaperones are proteins, which are often large and require an energy source such as ATP to function. A key molecular chaperone in *Escherichia coli* is GroEL, which consists of 14 subunits each of some 57.5 kD molecular mass arranged in two seven membered rings. There is a large cavity in the GroEL ring system, and it is widely believed that the cavity is required for successful  
30 protein folding activity. For optimal activity, a co-chaperone, GroES, is required which consists of a seven membered ring of 10 kD subunits. The activity of the GroEL/GroES complex requires energy source ATP.

Some proteins are monomers, consisting of a single subunit. Many proteins are oligomeric, consisting of more than one subunit. Sometimes the subunits are identical, sometimes there are different types of subunits. Frequently, the subunits are linked non-covalently. Sometimes, the subunits are attached covalently, with a stretch of polypeptide linking the C-terminus of one domain to the N-terminus of another.

Allosteric proteins are a special class of oligomeric proteins, which alternate between two or more different three-dimensional structures on the binding of ligands and substrates. Allosteric proteins are often involved in control processes in biology or where mechanical and physico-chemical energies are interconverted.

GroEL is an allosteric protein. The role of ATP is to trigger this allosteric change, causing GroEL to convert from a state that binds denatured proteins tightly to one that binds denatured proteins weakly. The co-chaperone, GroES, aids in this process by favouring the weak-binding state. It may also act as a cap, sealing off the cavity of GroEL. Further, its binding to GroEL is likely directly to compete with the binding of denatured substrates. The net result is that the binding of GroES and ATP to GroEL which has a substrate bound in its denatured form is to release the denatured substrate either into the cavity or into solution where it can refold.

Minichaperones have been described in detail elsewhere (see International patent application WO99/05163, the disclosure of which is incorporated herein by reference). Minichaperone polypeptides possess chaperoning activity when in monomeric form and do not require energy in the form of ATP. Defined fragments of the apical domain of GroEL of approximately 143-186 amino acid residues in length have molecular chaperone activity towards proteins either in solution under monomeric conditions or when monodisperse and attached to a support.

The activity of minichaperones, although sufficient for many purposes, is inferior to that of intact GroEL. There is thus a need for a more active form of minichaperone, which nevertheless retains independence from energy requirements.

## Summary of the Invention

The activity of minichaperones, although sufficient for many purposes, is inferior to that of intact GroEL. It is postulated that this could be due to the inability of GroES to oligomerise. There is thus a widespread requirement for a system which would allow the oligomerisation of polypeptides to form functional protein oligomers which have activities which surpass those of recombinant monomeric polypeptides.

According to the present invention, there is provided a polypeptide monomer capable of oligomerisation, said monomer comprising a polypeptide sequence which potentiates protein folding inserted into the sequence of a subunit of an oligomerisable protein scaffold.

It has been observed that the oligomerisation of polypeptide sequences which potentiate protein folding polypeptides allows their spatial juxtaposition, which potentiates their activity.

A "polypeptide sequence which potentiates protein folding" may any one of a number of polypeptides which are capable of facilitating the correct folding or refolding of proteins or polypeptides *in vivo* and/or *in vitro*. For example, the polypeptide sequence which potentiates protein folding may be a minichaperone polypeptide or a foldase. Advantageously, the foldase is selected from the group consisting of thiol/disulphide oxidoreductases and peptidyl prolyl isomerases. Preferably, the thiol/disulphide oxidoreductase is selected from the group consisting of *E. coli* DsbA and mammalian PDI, or a derivative thereof. Preferably, the peptidyl prolyl isomerase is a cyclophilin.

In a preferred aspect, the polypeptide sequence which potentiates protein folding may be a protease prosequence. Whereas molecular chaperones do not provide steric information necessary for newly synthesised proteins to fold correctly, but minimise the risk that partially folded protein chains will aggregate with one another, protease prosequences, which are intramolecular chaperones, provide steric information indispensable for the rest of the protein to fold correctly.

- A "minichaperone polypeptide", as referred to herein, refers to a minichaperone polypeptide as described in WO99/05163, the disclosure of which is incorporated herein by reference. The minichaperone polypeptide preferably comprises fragments of a molecular chaperone, preferably fragments of any hsp-60 chaperone, and may be selected from the group consisting of mammalian hsp-60 and GroEL, or a derivative thereof.

- In the case that the fragment is a fragment of GroEL, it advantageously does not have an Alanine residue at position 262 and/or an Isoleucine residue at position 267 of the sequence of intact GroEL as defined at GenBank Accession No. P06159. Preferably, it has a Leucine residue at position 262 and/or a Methionine residue at position 267 of the sequence of intact GroEL. The invention therefore encompasses the use of a fragment of GroEL comprising a Leucine residue at position 262 and/or a Methionine residue at position 267 of the sequence of intact GroEL for potentiating the folding of a polypeptide.
- In a preferred embodiment, the minichaperone comprises a region which is homologous to at least one of fragments 191-376, 191-345 and 191-335 of the sequence of intact GroEL.

- A protein scaffold is a protein, or part thereof, whose function is to determine the structure of the protein itself, or of a group of associated proteins or other molecules. Scaffolds therefore have a defined three-dimensional structure when assembled, and have the capacity to support molecules or polypeptide domains in or on the said structure. Advantageously, a scaffold has the ability to assume a variety of viable geometries, in relation to the three-dimensional structure of the scaffold and/or the insertion site of the polypeptides.

- Preferably, the scaffold according to the invention is a chaperonin cpn10/Hsp10 scaffold. Cpn10 is a widespread component of the cpn60/cpn10 chaperonin system. Examples of cpn10 include bacterial GroES and bacteriophage T4 Gp31. Further members of the cpn10 family will be known to those skilled in the art.

The invention moreover comprises the use of derivatives of naturally-occurring scaffolds. Derivatives of scaffolds (including scaffolds of the cpn10 and 60 families) comprise

mutants thereof, which may contain amino acid deletions, additions or substitutions (especially replacement of Cys residues in Gp31), hybrids formed by fusion of different members of the Cpn10 or 60 families and/or circularly permuted protein scaffolds, subject to the maintenance of the "oligomerisation" property described herein.

5

Protein scaffold subunits assemble to form a protein scaffold. In the context of the present invention, the scaffold may have any shape and may comprise any number of subunits. Preferably, the scaffold comprises between 2 and 20 subunits, advantageously between 5 and 15 subunits, and ideally about 10 subunits. The scaffold of cpn10 family members comprises seven subunits, in the shape of a seven-membered ring or annulus. Advantageously, therefore, the scaffold is a seven-membered ring.

10

15

Advantageously, the polypeptide sequence which potentiates protein folding is inserted into the sequence of the oligomerisable protein scaffold subunit such that both the N and C termini of the polypeptide monomer are formed by the sequence of the oligomerisable protein scaffold subunit. Thus, the polypeptide is included with the sequence of the scaffold subunit, for example by replacing one or more amino acids thereof.

20

25

It is known that cpn10 subunits possess a "mobile loop" within their structure. The mobile loop is positioned between amino acids 15 and 34, preferably between amino acids 16 to 33, of the sequence of *E. coli* GroES, and equivalent positions on other members of the cpn10 family. The mobile loop of T4 Gp31 is located between residues 22 to 45, advantageously 23 to 44. Advantageously, the polypeptide sequence which potentiates protein folding is inserted by replacing all or part of the mobile loop of a cpn10 family polypeptide.

30

Where the protein scaffold subunit is a cpn10 family polypeptide, the polypeptide sequence which potentiates protein folding may moreover be incorporated at the N or C terminus thereof, or in positions which are equivalent to the roof  $\beta$  hairpin of cpn10 family peptides. This position is located between positions 54 and 67, advantageously 55 to 66, and preferably 59 and 61 of bacteriophage T4 Gp31, or between positions 43 to 63, preferably 44 to 62, advantageously 56 to 57 of *E. coli* GroES.

Advantageously, the polypeptide sequence which potentiates protein folding may be inserted at an N or C terminus of a scaffold subunit in association with circular permutation of the subunit itself. Circular permutation is described in Graf and Schachman, PNAS(USA) 1996, 93:11591. Essentially, the polypeptide is circularised by fusion of the existing N and C termini, and cleavage of the polypeptide chain elsewhere to create novel N and C termini. In a preferred embodiment of the invention, the polypeptide may be included at the N and/or C terminus formed after circular permutation. The site of formation of the novel termini may be selected according to the features desired, and may include the mobile loop and/or the roof  $\beta$  hairpin.

10

Advantageously, polypeptide sequences which potentiates protein folding, which may be the same or different, may be inserted at more than one of the positions above-identified in the protein scaffold subunit. Thus, each subunit may comprise two or more polypeptides, which are displayed on the scaffold when this is assembled.

15

Polypeptide sequences which potentiate protein folding may be displayed on a scaffold subunit in free or constrained form, depending on the degree of freedom provided by the site of insertion into the scaffold sequence. For example, varying the length of the sequences flanking the mobile loop in the scaffold will modulate the degree of constraint of any polypeptide inserted therein.

20

In a second aspect, the invention relates to a polypeptide oligomer comprising two or more monomers according to the first aspect of the invention. The oligomer may be configured as a heterooligomer, comprising two or more different polypeptide sequences which potentiate protein folding inserted into the scaffold, or as a homooligomer, in which the polypeptides inserted into the scaffold are the same.

25

If the oligomer according to the invention is a heterooligomer, it may be configured such that the polypeptides juxtaposed thereon have complementary biological activities. For example, one or more minichaperones and foldases are advantageously displayed on the same scaffold, enabling them to act in concert.

30

10007344.10001

The monomers which constitute the oligomer may be covalently crosslinked to each other. Cross linking may be performed by recombinant approaches, such that the monomers are expressed *ab initio* as an oligomer; alternatively, cross-linking may be performed at Cys residues in the scaffold. For example, unique Cys residues inserted  
5 between positions 56 and 57 of the GroES scaffold, or equivalent positions on other members of the cpn10 family, may be used to cross-link scaffold subunits.

In a third aspect, the present invention relates to a method for preparing a polypeptide monomer capable of oligomerisation according to the first aspect of the invention,  
10 comprising the steps of inserting a nucleic acid sequence encoding a polypeptide sequence which potentiates protein folding into a nucleic acid sequence encoding a subunit of an oligomerisable protein scaffold, incorporating the resulting nucleic acid into an expression vector, and expressing the nucleic acid to produce the polypeptide monomers.

Oligomerised protease prosequences form polyvalent steric chaperones. The invention therefore provides polyvalent prosequence polypeptides, optionally incorporating one or more minichaperones and/or foldases, which are useful in refolding polypeptides *in vitro* and *in vivo*. Oligomerised prosequences may be used to potentiate the folding of, *inter*  
15 *alia*, proteases. Moreover, polyvalent prosequence polypeptides may be used to alter the folding patterns of polypeptides, thus permanently altering their activities.

The invention moreover relates to a method for producing a polypeptide oligomer according to the second aspect of the invention, comprising allowing the polypeptide  
20 monomers produced as above to associate into an oligomer. Preferably, the monomers are cross-linked to form the oligomer.

According to a fourth aspect of the present invention, there is provided a method for potentiating the folding of a polypeptide comprising contacting the polypeptide with a  
30 multimeric minichaperone polypeptide as described above.

The polypeptide which is folded by the method of the invention is preferably an unfolded or misfolded polypeptide, and advantageously comprises a disulphide.



The invention moreover concerns a method as described above wherein the oligomer is immobilised onto a solid phase support, which may be agarose. Accordingly, the invention also provides a solid phase support having immobilised thereon an oligomer according to the present invention, and a column packed at least in part with such a solid phase support.

The oligomer according to the invention may be combined with different oligomers, or with independent molecules capable of potentiating protein folding. For example, the oligomers according to the invention may be used, in solution or in immobilised form, in combination with foldases, chaperones or other enzymes.

### Brief Description of the Drawings

**Figure 1.** (a) Three-dimensional structure of Gp31 of bacteriophage T4 solved at 2.3 Å. Positions mentioned in the text are indicated (residues numbered as in van der Vies, S., Gatenby, A. & Georgopoulos, C. (1994) Nature 368, 654-656). (b) Three-dimensional structure of minichaperone GroEL(191-376) solved at 1.7 Å. The distance between residues 25 and 43 of Gp31 is around 12 Å; the distance between residues 191 and 376 of GroEL is around 9 Å. Positions mentioned in the text are indicated (residues numbered as in Hemmingsen, S. M., Woolford, C., van, d. V. S., Tilly, K., Dennis, D. T., Georgopoulos, C. P., Hendrix, R. W. & Ellis, R. J. (1988) Nature 333, 330-334). Secondary structure representations are drawn with MolScript (Kraulis, P. (1991) J. Appl. Crystallogr. 24, 946-950).

**Figure 2.** Schematic representation of Gp31 proteins in the vectors used in this study. The presence of the Gp31 mobile loop (residues 23 to 44) and/or minichaperone GroEL (residues 191 to 376) are indicated by boxes. The nucleotide sequence of the Gp31 mobile loop and relevant restriction sites are shown. The names of the corresponding vector are listed in the left margin.

**Figure 3.** (a) Molecular weight determination by analytical gel filtration chromatography. Wild-type proteins Gp31 ( $M_r \approx 7 \times 12$  kDa) and GroEL(191-376) ( $M_r \approx 22$  kDa) and,

Gp31Δloop and Gp31Δ::GroEL(191-376) (MC<sub>7</sub>) were run on a Superdex™ 200 (HR 10/30) column (Pharmacia Biotech.) calibrated with molecular weight standards (solid-line and circles). Gp31Δloop and MC<sub>7</sub> eluted at volumes corresponding to molecular weights of ≈145.6 and ≈215 kDa, respectively. (b) Molecular weight determination of MC<sub>7</sub> by equilibrium analytical ultracentrifugation. The apparent molecular weight of MC<sub>7</sub> is ≈215 kDa.

**Figure 4.** Characterisation of MC<sub>7</sub> by CD spectroscopy. (a) Far UV-CD spectrum at 25 °C. (b) Thermal denaturation followed at 222 nm at a heating rate of 1 °C.min<sup>-1</sup>.

**Figure 5.** (a) Binding specificity of MC<sub>7</sub> to GroES determined by ELISA. (b) Inhibition of MC<sub>7</sub> binding to heptameric co-chaperonin GroES by varying concentrations of synthetic peptide corresponding to residues 16 to 32 of GroES mobile loop determined by competition ELISA.

**Figure 6.** Binding avidity of MC<sub>7</sub> to anti-GroEL antibodies determined (a) by direct ELISA or (b) by indirect ELISA.

**Figure 7.** *In vitro* refolding of heat- and dithiothreitol-denatured mtMDH. (a) Protection of aggregation at 47 °C followed by light scattering at 550 nm. (b) Time-dependent reactivation of mtMDH at 25 °C. (c) Yields of mtMDH reactivation.

**Figure 8 and Figure 9** show the possible insertion sites for polypeptides in bacteriophage T4 Gp31.

**Figure 10** illustrates the potential attachment sites for polypeptides to a circular scaffold, in this case Gp31.

## Detailed Description of the Invention

### Definitions

- 5 *Oligomerisable scaffold.* An oligomerisable scaffold, as referred to herein, is a polypeptide which is capable of oligomerising to form a scaffold and to which a polypeptide may be fused, preferably covalently, without abolishing the oligomerisation capabilities. Thus, it provides a "scaffold" using which polypeptides may be arranged into multimers in accordance with the present invention. Optionally, parts of the wild-  
10 type polypeptide from which the scaffold is derived may be removed, for example by replacement with the polypeptide which is to be presented on the scaffold.

- Monomer.* Monomers according to the present invention are polypeptides which possess the potential to oligomerise. This is brought about by the incorporation, in the  
15 polypeptide, of an oligomerisable scaffold subunit which will oligomerise with further scaffold subunits if combined therewith.

- Oligomer.* As used herein, "oligomer" is synonymous with "polymer" or "multimer" and is used to indicate that the object in question is not monomeric. Thus, oligomeric  
20 polypeptides according to the invention comprise at least two monomeric units joined together covalently or non-covalently. The number of monomeric units employed will depend on the intended use of the oligomer, and may be between 2 and 20 or more. Advantageously, it is between 5 and 10, and preferably about 7.

- 25 *Polypeptide.* As used herein, a polypeptide is a molecule comprising at least one peptide bond linking two amino acids. This term is synonymous with "protein" and "peptide", both of which are used in the art to describe such molecules. A polypeptide may comprise other, non-amino acid components. The polypeptide the folding of which is potentiated by the method of the invention may be any polypeptide. Preferably, however,  
30 it is an unfolded or misfolded polypeptide which is in need of folding. Alternatively, however, it may be a folded polypeptide which is to be maintained in a folded state (see below).

10007214.110801

Preferably, the polypeptide which is folded according to the invention contains at least one disulphide. Such polypeptides may be referred to herein as *disulphide-containing polypeptides*.

- 5 Examples of polypeptides include those used for medical or biotechnological use, such as interleukins, interferons, antibodies and their fragments, insulin, transforming growth factor, and many toxins and proteases.

- 10 *Potentiating the folding.* The invention envisages at least two situations. A first situation is one in which the polypeptide to be folded is in an unfolded or misfolded state, or both. In this case, its correct folding is potentiated by the method of the invention. A second situation is one in which the polypeptide is substantially already in its correctly folded state, that is all or most of it is folded correctly or nearly correctly. In this case, the method of the invention serves to maintain the folded state of the polypeptide by affecting
- 15 the folded/unfolded equilibrium so as to favour the folded state. This prevents loss of activity of an already substantially correctly folded polypeptide. These, and other, eventualities are covered by the reference to "potentiating" the folding of the polypeptide.

- 20 *Contacting.* The reagents used in the method of the invention require physical contact with the polypeptides whose folding is to be potentiated. This contact may occur in free solution, *in vitro* or *in vivo*, with one or more components of the reaction immobilised on solid supports. In a preferred aspect, the contact occurs with the minichaperone oligomer and/or the thiol/disulphide oxidoreductase immobilised on a solid support, for example on
- 25 which may be added to a solution comprising a polypeptide whose folding is to be potentiated.

- 30 *Fragment.* When applied to chaperone molecules, a fragment is anything other than the entire native molecular chaperone molecule which nevertheless retains chaperonin activity. Advantageously, a fragment of a chaperonin molecule remains monomeric in solution. Preferred fragments are described below. Advantageously, chaperone fragments are between 50 and 200 amino acids in length, preferably between 100 and 200 amino acids in length and most preferably about 150 amino acids in length. Fragments of

chaperone molecules which remain monomeric in solution and possess a chaperoning activity which is not energy-dependent are referred to as minichaperones.

5 *Unfolded.* As used herein, a polypeptide may be unfolded when at least part of it has not yet acquired is correct or desired secondary or tertiary structure. A polypeptide is *misfolded* when it has acquired an at least partially incorrect or undesired secondary or tertiary structure.

10 *Immobilised, immobilising.* Permanently attached, covalently or otherwise. In a preferred aspect of the present invention, the term "immobilise", and grammatical variations thereof, refer to the attachment of molecular chaperones or, preferably, foldase polypeptides to a solid phase support using a method as described in WO99/05163.

15 *Solid (phase) support.* Reagents used in the invention may be immobilised onto solid phase supports. This means that they are permanently attached to an entity which remains in a different (solid) phase from reagents which are in solution. For example, the solid phase could be in the form of beads, a "polypeptide chip", a resin, a matrix, a gel, the material forming the walls of a vessel or the like. Matrices, and in particular gels, such as agarose gels, may conveniently be packed into columns. A particular advantage  
20 of solid phase immobilisation is that the reagents may be removed from contact with the polypeptide(s) with facility.

25 *Foldase.* In general terms, a foldase is an enzyme which participates in the potentiation of protein folding through its enzymatic activity to catalyse the rearrangement or isomerisation of bonds in the folding polypeptide. They are thus distinct from a molecular chaperone, which bind to polypeptides in unstable or non-native structural states and potentiate correct folding without enzymatic catalysis of bond rearrangement. Many classes of foldase are known, and they are common to animals, plants and bacteria. They include peptidyl prolyl isomerases and thiol/disulphide oxidoreductases. The invention  
30 comprises the use of all foldases which are capable of potentiating protein folding through covalent bond rearrangement.

Moreover, as used herein, the term "a foldase" includes one or more foldases. In general, in the present specification the use of the singular does not preclude the presence of a plurality of the entities referred to, unless the context specifically requires otherwise.

5 *Thiol/disulphide oxidoreductase*. As the name implies, thiol/disulphide oxidoreductases catalyse the formation of disulphide bonds and can thus dictate the folding rate of disulphide-containing polypeptides. The invention accordingly comprises the use of any polypeptide possessing such an activity. This includes chaperone polypeptides, or fragments thereof, which may possess PDI activity (Wang & Tsou, (1998) FEBS lett.  
10 425:382-384). In Eukaryotes, thiol/disulphide oxidoreductases are generally referred to as PDIs (protein disulphide isomerases). PDI interacts directly with newly synthesised secretory proteins and is required for the folding of nascent polypeptides in the endoplasmic reticulum (ER) of eukaryotic cells. Enzymes found in the ER with PDI activity include mammalian PDI (Edman *et al.*, 1985, Nature 317:267, yeast PDI  
15 (Mizunaga *et al.* 1990, J. Biochem. 108:848), mammalian Erp59 (Mazzarella *et al.*, 1990, J. Biochem. 265:1094), mammalian prolyl-4-hydroxylase (Pihlajaniemi *et al.* , 1987, EMBO J. 6: 643) yeast GSBP (Lamantia *et al.*, 1991, Proc. Natl. Acad. Sci. USA, 88:4453) and mammalian T3BP (Yamauchi *et al.*, 1987, Biochem. Biophys. Res. Commun. 146:1485), *A. niger* PdiA (Ngiam *et al.*, (1997) Curr. genet. 31:133-138) and  
20 yeast EUGI (Tachibana *et al.*, 1992, Mol. Cell Biol. 12, 4601). In prokaryotes, equivalent proteins exist, such as the DsbA protein of *E. coli*. Other peptides with similar activity include, for example, p52 from *T. cruzi* (Moutiez *et al.*, (1997) Biochem. J. 322:43-48). These polypeptides, and other functionally equivalent polypeptides, are included with the scope of the present invention, as are derivatives of the polypeptides which share the  
25 relevant activity (see below). Preferably, the thiol/disulphide oxidoreductase according to the invention is selected from the group consisting of mammalian PDI or *E. coli* DsbA.

*Peptidyl-prolyl isomerase*. Peptidyl-prolyl isomerases are known enzymes widely present in a variety of cells. Examples include cyclophilin (see, for example, Bergsma  
30 *et al.* (1991) J. Biol. Chem. 266:23204-23214), parbullen, SurA (Rouviere and Gross, (1996) Genes Dev. 10:3170-3182) and FK506 binding proteins FKBP51 and FKBP52. PPI is responsible for the *cis-trans* isomerisation of peptidyl-prolyl bonds in polypeptides, thus potentiating correct folding. The invention includes any polypeptide having PPI activity.

This includes chaperone polypeptides, or fragments thereof, which may possess PPI activity (Wang & Tsou, (1998) FEBS lett. 425:382-384).

- 5 *Molecular Chaperone.* Chaperones, or chaperonins, are polypeptides which potentiate protein folding by non-enzymatic means, in that they do not catalyse the chemical modification of any structures in folding polypeptides, by potentiate the correct folding of polypeptides by facilitating correct structural alignment thereof. Molecular chaperones are well known in the art, several families thereof being characterised. The invention is applicable to any molecular chaperone molecule, which term includes, for example, the
- 10 molecular chaperones selected from the following non-exhaustive group:

p90 Calnexin	Salopek <i>et al.</i> , J. Investig Dermatol Symp Proc (1996) 1:195
HSP family	Walsh <i>et al.</i> , Cell Mol. Life Sci. (1997) 53:198
HSP 70 family	Rokutan <i>et al.</i> , J. Med. Invest. (1998) 44:137
DNA K	Rudiger <i>et al.</i> , Nat. Struct. Biol. (1997) 4:342
DNAJ	Cheetham <i>et al.</i> , Cell Stress Chaperones (1998) 3:28
HSP 60 family; GroEL	Richardson <i>et al.</i> , Trends Biochem. (1998) 23:138
ER-associated chaperones	Kim <i>et al.</i> , Endocr Rev (1998) 19:173
HSP 90	Smith, Biol. Chem. (1988) 379:283
Hsc 70	Hohfeld, Biol. Chem. (1988) 379:269
sHsps; SecA; SecB	Beissinger <i>et al.</i> , Biol. Chem. (1988) 379:245
Trigger factor	Wang <i>et al.</i> , FEBS Lett. (1998) 425:382
zebrafish hsp 47, 70 and 90	Krone <i>et al.</i> , Biochem. Cell Biol. (1997) 75:487
HSP 47	Nagata, Matrix Biol. (1998) 16:379
GRP 94	Nicchitta <i>et al.</i> , Curr. Opin. Immunol. (1998) 10:103
Cpn 10	Cavanagh, Rev. Reprod. (1996) 1:28
BiP	Sommer <i>et al.</i> , FASEB J. (1997) 11:1227
GRP 78	Brostrom <i>et al.</i> , Prog. Nucl. Acid. res. Mol. Biol. (1998) 58:79
C1p, FtsH	Suzuki <i>et al.</i> , Trends Biochem. Sci. (1997) 22:118
Ig invariant chain	Weenink <i>et al.</i> Immunol. Cell biol. (1997) 75:69
mitochondrial hsp 70	Horst <i>et al.</i> , BBA (1997) 1318:71

EBP	Hinek, Arch. Immunol. Ther. Exp. (1997) 45:15
mitochondrial m-AAA	Langer <i>et al.</i> , Experientia (1996) 52:1069
Yeast Ydj1	Lyman <i>et al.</i> , Experientia (1996) 52:1042
Hsp 104	Tuite <i>et al.</i> , Trends Genet. (1996) 12:467
ApoE	Blain <i>et al.</i> , Presse Med. (1996) 25:763
Syc	Wattiau <i>et al.</i> , Mol. Microbiol. (1996) 20:255
Hip	Ziegelhoffer <i>et al.</i> , Curr. Biol. (1996) 6:272
TriC family	Hendrick <i>et al.</i> , FASEB J. (1995) 9:1559
CCT	Kubota <i>et al.</i> , Eur. J. Biochem. (1995) 230:3
PapD, calmodulin	Stanfield <i>et al.</i> , Curr. Opin. Struct. Biol. (1995) 5:103

Two major families of protein folding chaperones which have been identified, the heat shock protein 60 (hsp60) class and the heat shock protein 70 (hsp70) class, are especially preferred for use herein. Chaperones of the hsp-60 class are structurally distinct from chaperones of the hsp-70 class. In particular, hsp-60 chaperones appear to form a stable scaffold of two heptamer rings stacked one atop another which interacts with partially folded elements of secondary structure. On the other hand, hsp-70 chaperones are monomers or dimers and appear to interact with short extended regions of a polypeptide.

Hsp70 chaperones are well conserved in sequence and function. Analogues of hsp-70 include the eukaryotic hsp70 homologue originally identified as the IgG heavy chain binding protein (BiP). BiP is located in all eukaryotic cells within the lumen of the endoplasmic reticulum (ER). The prokaryotic DnaK hsp70 protein chaperone in *Escherichia coli* shares about 50% sequence homology with an hsp70 KAR2 chaperone in yeast (Rose *et al.* 1989 Cell 57:1211-1221). Moreover, the presence of mouse BiP in yeast can functionally replace a lost yeast KAR2 gene (Normington *et al.* 19: 1223-1236).

Hsp-60 chaperones are universally conserved (Zeilstra-Ryalls *et al.*, (1991) Ann. Rev. Microbiol. 45:301-325) and include hsp-60 homologues from large number of species, including man. They include, for example, the *E. coli* GroEL polypeptide; *Ehrlichia sennetsu* GroEL (Zhang *et al.*, (1997) FEMS Immunol. Med. Microbiol. 18:39-46); *Trichomonas vaginalis* hsp-60 (Bozner *et al.*, (1997) J. Parasitol. 83:224-229; rat hsp-60



(Venner *et al.*, (1990) NAR 18:5309; and yeast hsp-60 (Johnson *et al.*, (1989) Gene 84:295-302.

In a preferred aspect, the present invention relates to fragments of polypeptides of the hsp-60 family. These proteins being universally conserved, any member of the family may be used; however, in a particularly advantageous embodiment, fragments of GroEL, such as *E. coli* GroEL, are employed. It has also found that agarose-immobilised calmodulin does have a chaperoning activity, presumably because of its exposed hydrophobic groups.

The sequence of GroEL is available in the art and from academic databases (see GenBank Accession No. P06159); however, GroEL fragments which conform to the database sequence are inoperative. Specifically, the database contains a sequence in which positions 262 and 267 are occupied by Alanine and Isoleucine respectively. Fragments incorporating one or both of these residues at these positions are inoperative and unable to potentiate the folding of polypeptides. The invention, instead, relates to a GroEL polypeptide in which at least one of positions 262 and 267 is occupied by Leucine and Methionine respectively.

*Derivative.* The present invention relates to derivatives of molecular chaperones, peptidyl-prolyl isomerases and thiol/disulphide oxidoreductases. In a preferred aspect, therefore, the terms "molecular chaperone", "peptidyl-prolyl isomerase" and "thiol-disulphide oxidoreductase" include derivatives thereof which retain the stated activity. The derivatives provided by the present invention include splice variants encoded by mRNA generated by alternative splicing of a primary transcript, amino acid mutants, glycosylation variants and other covalent derivatives of molecular chaperones or foldases which retain the functional properties of molecular chaperones, peptidyl-prolyl isomerases and/or thiol/disulphide oxidoreductases. Exemplary derivatives include molecules which are covalently modified by substitution, chemical, enzymatic, or other appropriate means with a moiety other than a naturally occurring amino acid. Such a moiety may be a detectable moiety such as an enzyme or a radioisotope. Further included are naturally occurring variants of molecular chaperones or foldases found within a particular species, whether mammalian, other vertebrate, yeast, prokaryotic or otherwise.

Such a variant may be encoded by a related gene of the same gene family, by an allelic variant of a particular gene, or represent an alternative splicing variant of a molecular chaperone or foldase. Possible derivatives of the polypeptides employed in the invention are described below.

5

## Description of Preferred Embodiments

### Scaffold Proteins

- 10 In a preferred embodiment, the scaffold polypeptide is based on members of the cpn10/Hsp10 family, such as GroES or an analogue thereof. A highly preferred analogue is the T4 polypeptide Gp31. GroES analogues, including Gp31, possess a mobile loop (Hunt, J. F., *et al.*, (1997) *Cell* **90**, 361-371; Landry, S. J., *et al.*, (1996) *Proc. Natl. Acad. Sci. U.S.A.* **93**, 11622-11627) which may be inserted into, or replaced, in order to fuse the polypeptide to the scaffold.

15

Cpn10 homologues are widespread throughout animals, plants and bacteria. For example, a search of GenBank indicates that cpn10 homologues are known in the following species:

20

- Actinobacillus actinomycetemcomitans*; *Actinobacillus pleuropneumoniae*; *Aeromonas salmonicida*; *Agrobacterium tumefaciens*; *Allochrodatum vinosum*; *Amoeba proteus symbiotic bacterium*; *Aquifex aeolicus*; *Arabidopsis thaliana*; *Bacillus sp*; *Bacillus stearothermophilus*; *Bacillus subtilis*; *Bartonella henselae*; *Bordetella pertussis*; *Borrelia burgdorferi*; *Brucella abortus*; *Buchnera aphidicola*; *Burkholderia cepacia*; *Burkholderia vietnamiensis*; *Campylobacter jejuni*; *Caulobacter crescentus*; *Chlamydia muridarum*; *Chlamydia trachomatis*; *Chlamydomydia pneumoniae*; *Clostridium acetobutylicum*; *Clostridium perfringens*; *Clostridium thermocellum*; *coliphage T*; *Cowdria ruminantium*; *Cyanella Cyanophora paradoxa*; *Ehrlichia canis*; *Ehrlichia chaffeensis*; *Ehrlichia equi*; *Ehrlichia phagocytophila*; *Ehrlichia risticii*; *Ehrlichia sennetsu*; *Ehrlichia sp 'HGE agent'*; *Enterobacter aerogenes*; *Enterobacter agglomerans*; *Enterobacter amnigenus*; *Enterobacter asburiae*; *Enterobacter gergoviae*; *Enterobacter intermedius*; *Erwinia aphidicola*; *Erwinia carotovora*; *Erwinia herbicola*; *Escherichia*

30

10007214.110801

*coli*; *Francisella tularensis*; *Glycine max*; *Haemophilus ducreyi*; *Haemophilus influenzae* Rd; *Helicobacter pylori*; *Holospira obtusa*; *Homo sapiens*; *Klebsiella ornithinolytica*; *Klebsiella oxytoca*; *Klebsiella planticola*; *Klebsiella pneumoniae*; *Lactobacillus helveticus*; *Lactobacillus zeae*; *Lactococcus lactis*; *Lawsonia intracellularis*; *Leptospira interrogans*; *Methylovorus* sp strain SS; *Mycobacterium avium*; *Mycobacterium avium* subsp *avium*; *Mycobacterium avium* subsp *paratuberculosis*; *Mycobacterium leprae*; *Mycobacterium tuberculosis*; *Mycoplasma genitalium*; *Mycoplasma pneumoniae*; *Myzus persicae* primary endosymbiont; *Neisseria gonorrhoeae*; *Oscillatoria* sp NKBG; *Pantoea ananas*; *Pasteurella multocida*; *Porphyromonas gingivalis*; *Pseudomonas aeruginosa*;

10 *Pseudomonas aeruginosa*; *Pseudomonas putida*; *Rattus norvegicus*; *Rattus norvegicus*; *Rhizobium leguminosarum*; *Rhodobacter capsulatus*; *Rhodobacter sphaeroides*; *Rhodothermus marinus*; *Rickettsia prowazekii*; *Rickettsia rickettsii*; *Saccharomyces cerevisiae*; *Serratia ficaria*; *Serratia marcescens*; *Serratia rubidaea*; *Sinorhizobium meliloti*; *Sitophilus oryzae* principal endosymbiont; *Stenotrophomonas maltophilia*;

15 *Streptococcus pneumoniae*; *Streptomyces albus*; *Streptomyces coelicolor*; *Streptomyces coelicolor*; *Streptomyces lividans*; *Synechococcus* sp; *Synechococcus vulcanus*; *Synechocystis* sp; *Thermoanaerobacter brockii*; *Thermotoga maritima*; *Thermus aquaticus*; *Treponema pallidum*; *Wolbachia* sp; *Zymomonas mobilis*.

20 An advantage of cpn10 family subunits is that they possess a mobile loop, responsible for the protein folding activity of the natural chaperonin, which may be removed without affecting the scaffold.

Cpn10 with a deleted mobile loop possesses no biological activity, making it an advantageously inert scaffold, thus minimising any potentially deleterious effects. Insertion of an appropriate biologically active polypeptide can confer a biological activity on the novel polypeptide thus generated. Indeed, the biological activity of the inserted polypeptide may be improved by incorporation of the biologically active polypeptide into the scaffold.

30 Alternative sites for peptide insertion are possible. An advantageous option is in the position equivalent to the roof beta hairpin in GroES. This involves replacement of Glu-60 in Gp31 by the desired peptide. The amino acid sequence is Pro(59)-Glu(60)-Gly(61).

This is conveniently converted to a SmaI site at the DNA level (CCC:GGG) encoding Pro-Gly, leaving a blunt-ended restriction site for peptide insertion as a DNA fragment. Similarly, an insertion may be made at between positions 56 and 57 of the GroES sequence, and at equivalent positions in other cpn10 family members. Alternatively,  
5 inverse PCR may be used, to display the peptide on the opposite side of the scaffold.

Members of the cpn60/Hsp60 family of chaperonin molecules may also be used as scaffolds. For example, the tetradecameric bacterial chaperonin GroEL may be used. Advantageously, polypeptides would be inserted between positions 191 and 376, in  
10 particular between positions 197 and 333 (represented by *SacII* engineered and unique *Cla I* sites) to maintain intact the hinge region between the equatorial and the apical domains in order to impart mobility to the inserted polypeptide. The choice of scaffold may depend upon the intended application of the oligomer: for example, if the oligomer is intended for vaccination purposes (see below), the use of an immunogenic scaffold, such  
15 as that derived from *Mycobacterium tuberculosis*, is highly advantageous and confers an adjuvant effect.

Mutants of cpn60 molecules may also be used. For example, the single ring mutant of GroEL (GroELSR1) contains four point mutations which effect the major attachment  
20 between the two rings of GroEL (R452E, E461A, S463A and V464A) and is functionally inactive *in vitro* because it is release to bind GroES. GroELSR2 has an additional mutation at Glu191-Gly, which restores activity by reducing the affinity for GroES. Both of these mutants for ring structures and would be suitable for use as scaffolds.

## 25 Configurations of Oligomers according to the Invention

Figures 8 – 10 show various topologies and applications for scaffolded polypeptides in accordance with the present invention. In figures 8 and 9, the possible insertion sites for polypeptides are shown. Insertion of polypeptides may be performed by any suitable  
30 technique, including those set forth by Doi and Yanagawa (FEBS Letters (1999) 457:1-4). As set forth therein, insertion of polypeptides may be combined with randomisation to produce libraries of polypeptide repertoires, suitable for display and selection.

Figure 10 illustrates the potential attachment sites for polypeptides to a circular scaffold, in this case Gp31. Reading from left to right, the figure shows: no attachment, attachment to the mobile loop, attachment to the roof  $\beta$  hairpin, attachment at both the mobile loop and the roof  $\beta$  hairpin, attachment at the C terminus, attachment at both N and C termini, attachment at both N and C termini and the mobile loop, and attachment at both N and C termini, the roof  $\beta$  hairpin and the mobile loop. As will be apparent, further configurations are possible, and can be combined in any way in the heptamer, leading to a total of  $5.4 \times 10^8$  possible configurations.

#### Recombinant DNA techniques

The present invention advantageously makes use of recombinant DNA technology in order to construct polypeptide monomers and oligomers. Advantageously, polypeptide monomers or oligomers may be expressed from nucleic acid sequences which encode them.

As used herein, vector (or plasmid) refers to discrete elements that are used to introduce heterologous DNA into cells for either expression or replication thereof. Selection and use of such vehicles are well within the skill of the artisan. Many vectors are available, and selection of appropriate vector will depend on the intended use of the vector, i.e. whether it is to be used for DNA amplification or for DNA expression, the size of the DNA to be inserted into the vector, and the host cell to be transformed with the vector. Each vector contains various components depending on its function (amplification of DNA or expression of DNA) and the host cell for which it is compatible. The vector components generally include, but are not limited to, one or more of the following: an origin of replication, one or more marker genes, an enhancer element, a promoter, a transcription termination sequence and a signal sequence.

Both expression and cloning vectors generally contain nucleic acid sequence that enable the vector to replicate in one or more selected host cells. Typically in cloning vectors, this sequence is one that enables the vector to replicate independently of the host chromosomal DNA, and includes origins of replication or autonomously replicating sequences. Such sequences are well known for a variety of bacteria, yeast and viruses.

The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria, the 2m plasmid origin is suitable for yeast, and various viral origins (e.g. SV 40, polyoma, adenovirus) are useful for cloning vectors in mammalian cells. Generally, the origin of replication component is not needed for mammalian expression vectors unless  
5 these are used in mammalian cells competent for high level DNA replication, such as COS cells.

Most expression vectors are shuttle vectors, i.e. they are capable of replication in at least one class of organisms but can be transfected into another class of organisms for  
10 expression. For example, a vector is cloned in *E. coli* and then the same vector is transfected into yeast or mammalian cells even though it is not capable of replicating independently of the host cell chromosome. DNA can be amplified by PCR and be directly transfected into the host cells without any replication component.

Advantageously, an expression and cloning vector may contain a selection gene also referred to as selectable marker. This gene encodes a protein necessary for the survival or growth of transformed host cells grown in a selective culture medium. Host cells not transformed with the vector containing the selection gene will not survive in the culture medium. Typical selection genes encode proteins that confer resistance to antibiotics and  
15 other toxins, e.g. ampicillin, neomycin, methotrexate or tetracycline, complement auxotrophic deficiencies, or supply critical nutrients not available from complex media.

As to a selective gene marker appropriate for yeast, any marker gene can be used which facilitates the selection for transformants due to the phenotypic expression of the marker  
25 gene. Suitable markers for yeast are, for example, those conferring resistance to antibiotics G418, hygromycin or bleomycin, or provide for prototrophy in an auxotrophic yeast mutant, for example the URA3, LEU2, LYS2, TRP1, or HIS3 gene.

Since the replication of vectors is conveniently done in *E. coli*, an *E. coli* genetic marker  
30 and an *E. coli* origin of replication are advantageously included. These can be obtained from *E. coli* plasmids, such as pBR322, Bluescript<sup>®</sup> vector or a pUC plasmid, e.g. pUC18 or pUC19, which contain both *E. coli* replication origin and *E. coli* genetic marker conferring resistance to antibiotics, such as ampicillin.

Suitable selectable markers for mammalian cells are those that enable the identification of cells which have been transformed, such as dihydrofolate reductase (DHFR, methotrexate resistance), thymidine kinase, or genes conferring resistance to G418 or hygromycin. The mammalian cell transformants are placed under selection pressure which only those transformants which have taken up and are expressing the marker are uniquely adapted to survive. In the case of a DHFR or glutamine synthase (GS) marker, selection pressure can be imposed by culturing the transformants under conditions in which the pressure is progressively increased, thereby leading to amplification (at its chromosomal integration site) of both the selection gene and the linked DNA that encodes the polypeptide according to the invention. Amplification is the process by which genes in greater demand for the production of a protein critical for growth, together with closely associated genes which may encode a desired protein, are reiterated in tandem within the chromosomes of recombinant cells. Increased quantities of desired protein are usually synthesised from thus amplified DNA.

Expression and cloning vectors usually contain a promoter that is recognised by the host organism and is operably linked to the heterologous nucleic acid coding sequence. Such a promoter may be inducible or constitutive. The promoters are operably linked to the coding sequence by inserting the isolated promoter sequence into the vector. Many heterologous promoters may be used to direct amplification and/or expression of the coding sequence. The term "operably linked" refers to a juxtaposition wherein the components described are in a relationship permitting them to function in their intended manner. A control sequence "operably linked" to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under conditions compatible with the control sequences.

Promoters suitable for use with prokaryotic hosts include, for example, the  $\beta$ -lactamase and lactose promoter systems, alkaline phosphatase, the tryptophan (trp) promoter system and hybrid promoters such as the tac promoter. Their nucleotide sequences have been published, thereby enabling the skilled worker operably to ligate them to the coding sequence, using linkers or adapters to supply any required restriction sites. Promoters for

use in bacterial systems will also generally contain a Shine-Delgarno sequence operably linked to the coding sequence.

Preferred expression vectors are bacterial expression vectors which comprise a promoter of a bacteriophage such as phagex or T7 which is capable of functioning in the bacteria.

In one of the most widely used expression systems, the nucleic acid encoding the fusion protein may be transcribed from the vector by T7 RNA polymerase (Studier et al, Methods in Enzymol. 185; 60-89, 1990). In the *E. coli* BL21(DE3) host strain, used in conjunction with pET vectors, the T7 RNA polymerase is produced from the  $\lambda$ -lysogen DE3 in the host bacterium, and its expression is under the control of the IPTG inducible lac UV5 promoter. This system has been employed successfully for over-production of many proteins. Alternatively the polymerase gene may be introduced on a lambda phage by infection with an int- phage such as the CE6 phage which is commercially available (Novagen, Madison, USA). other vectors include vectors containing the lambda PL promoter such as PLEX (Invitrogen, NL) , vectors containing the trc promoters such as pTrcHisXpressTm (Invitrogen) or pTrc99 (Pharmacia Biotech, SE) , or vectors containing the tac promoter such as pKK223-3 (Pharmacia Biotech) or PMAL (new England Biolabs, MA, USA).

Moreover, the coding sequence according to the invention preferably includes a secretion sequence in order to facilitate secretion of the polypeptide from bacterial hosts, such that it will be produced as a soluble native peptide rather than in an inclusion body. The peptide may be recovered from the bacterial periplasmic space, or the culture medium, as appropriate.

Suitable promoting sequences for use with yeast hosts may be regulated or constitutive and are preferably derived from a highly expressed yeast gene, especially a *Saccharomyces cerevisiae* gene. Thus, the promoter of the TRP1 gene, the ADHI or ADHII gene, the acid phosphatase (PH05) gene, a promoter of the yeast mating pheromone genes coding for the a- or  $\alpha$ -factor or a promoter derived from a gene encoding a glycolytic enzyme such as the promoter of the enolase, glyceraldehyde-3-phosphate dehydrogenase (GAP), 3-phospho glycerate kinase (PGK), hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-



phosphoglycerate mutase, pyruvate kinase, triose phosphate isomerase, phosphoglucose isomerase or glucokinase genes, the *S. cerevisiae* GAL 4 gene, the *S. pombe* nmt 1 gene or a promoter from the TATA binding protein (TBP) gene can be used. Furthermore, it is possible to use hybrid promoters comprising upstream activation sequences (UAS) of one yeast gene and downstream promoter elements including a functional TATA box of another yeast gene, for example a hybrid promoter including the UAS(s) of the yeast PH05 gene and downstream promoter elements including a functional TATA box of the yeast GAP gene (PH05-GAP hybrid promoter). A suitable constitutive PH05 promoter is e.g. a shortened acid phosphatase PH05 promoter devoid of the upstream regulatory elements (UAS) such as the PH05 (-173) promoter element starting at nucleotide -173 and ending at nucleotide -9 of the PH05 gene.

Transcription from vectors in mammalian hosts may be controlled by promoters derived from the genomes of viruses such as polyoma virus, adenovirus, fowlpox virus, bovine papilloma virus, avian sarcoma virus, cytomegalovirus (CMV), a retrovirus and Simian Virus 40 (SV40), from heterologous mammalian promoters such as the actin promoter or a very strong promoter, e.g. a ribosomal protein promoter, provided such promoters are compatible with the host cell systems.

Transcription of a coding sequence by higher eukaryotes may be increased by inserting an enhancer sequence into the vector. Enhancers are relatively orientation and position independent. Many enhancer sequences are known from mammalian genes (e.g. elastase and globin). However, typically one will employ an enhancer from a eukaryotic cell virus. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270) and the CMV early promoter enhancer. The enhancer may be spliced into the vector at a position 5' or 3' to the coding sequence, but is preferably located at a site 5' from the promoter.

Advantageously, a eukaryotic expression vector may comprise a locus control region (LCR). LCRs are capable of directing high-level integration site independent expression of transgenes integrated into host cell chromatin, which is of importance especially where the coding sequence is to be expressed in the context of a permanently-transfected

eukaryotic cell line in which chromosomal integration of the vector has occurred, in vectors designed for gene therapy applications or in transgenic animals.

Eukaryotic expression vectors will also contain sequences necessary for the termination of transcription and for stabilising the mRNA. Such sequences are commonly available from the 5' and 3' untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA.

- 10 An expression vector includes any vector capable of expressing nucleic acids that are operatively linked with regulatory sequences, such as promoter regions, that are capable of expression of such DNAs. Thus, an expression vector refers to a recombinant DNA or RNA construct, such as a plasmid, a phage, recombinant virus or other vector, that upon introduction into an appropriate host cell, results in expression of the cloned DNA.
- 15 Appropriate expression vectors are well known to those with ordinary skill in the art and include those that are replicable in eukaryotic and/or prokaryotic cells and those that remain episomal or those which integrate into the host cell genome. For example, nucleic acids may be inserted into a vector suitable for expression of cDNAs in mammalian cells, e.g. a CMV enhancer-based vector such as pEVRF (Matthias, et al., (1989) NAR 17, 6418).
- 20

Construction of vectors according to the invention employs conventional ligation techniques. Isolated plasmids or DNA fragments are cleaved, tailored, and religated in the form desired to generate the plasmids required. If desired, analysis to confirm correct sequences in the constructed plasmids is performed in a known fashion. Suitable methods for constructing expression vectors, preparing in vitro transcripts, introducing DNA into host cells, and performing analyses for assessing expression and function are known to those skilled in the art. Gene presence, amplification and/or expression may be measured in a sample directly, for example, by conventional Southern blotting, Northern blotting to quantitate the transcription of mRNA, dot blotting (DNA or RNA analysis), or in situ hybridisation, using an appropriately labelled probe which may be based on a sequence provided herein. Those skilled in the art will readily envisage how these methods may be modified, if desired.

25

30

Description of Preferred Embodiments

5 In a preferred aspect, the present invention relates to a method for oligomerising polypeptides and novel oligomeric polypeptides producible thereby. By incorporating polypeptides into a scaffold which is capable of oligomerising, as described herein, oligomers may be produced wherein the chosen polypeptides are juxtaposed. The polypeptides selected for oligomerisation may be the same or different; thus, it is possible to produce homooligomers or heterooligomers.

10 Preferably, the oligomeric proteins of the invention are oligomeric minichaperone polypeptides. Oligomeric minichaperones have been shown to possess particularly advantageous properties.

15 The invention thus relates to the use of oligomeric molecular chaperone fragments, whether alone or in combination with other polypeptides, for potentiating the folding or refolding of polypeptides.

20 The present invention may be practised in a number of configurations, according to the required use to which the invention is to be put. In a first configuration, the invention relates to the use of minichaperone oligomers alone to potentiate the folding or refolding of polypeptides. This may be performed *in vivo* or *in vitro*, in solution or on a solid support. For example, oligomerised minichaperones may be immobilised onto resins and packed into columns for use in refolding polypeptides which are passed through the column. Methods for immobilising minichaperones are described in International patent application WO99/05163, incorporated herein by reference.

25 In an alternative configuration, minichaperone oligomers according to the invention may be expressed *in vivo* or administered to cells or organisms *in vivo* in order to potentiate protein folding therein.

30 In a second configuration, the invention provides a combination of a molecular chaperone and a thiol/disulphide oxidoreductase to facilitate protein folding. One or both of the

chaperones and thiol/disulphide oxidoreductases may be incorporated into the scaffold according to the invention. The combination of a molecular chaperone and a thiol/disulphide oxidoreductase provides a synergistic effect on protein folding which results in a greater quantity of active, correctly folded protein being produced than would be expected from a merely additive relationship. Advantageously, one or more of the components used to potentiate protein folding in accordance with the present invention is immobilised on a solid support. However, both molecular chaperones and thiol/disulphide oxidoreductases may be used in solution. They may be used in free solution, but also in suspension, for example bound to a matrix such as beads, for example Sepharose beads, or bound to solid surfaces which are in contact with solutions, such as the inside surfaces of bottles containing solutions, test tubes and the like.

In a third configuration, the invention relates to the use of a combination of a molecular chaperone and a thiol/disulphide oxidoreductase with a peptidyl prolyl isomerase. The peptidyl prolyl isomerase may be present either incorporated into the oligomeric scaffold, bound to a solid support, or in solution. Moreover, it may be bound to beads suspended in solution. The peptidyl prolyl isomerases may be used together with a molecular chaperone alone, with a thiol/disulphide oxidoreductase alone, or with both a molecular chaperone and a thiol/disulphide oxidoreductase. In the latter case, further synergistic effects are apparent over the additive effects which would be expected from the use of the three components together. In particular, an increase in the proportion of the folded protein which is recovered as monodisperse protein, as opposed to aggregated protein, increases substantially.

Used in accordance with any of the foregoing configurations, or otherwise in accordance with the following claims, the invention may be used to facilitate protein folding in a variety of situations. For example, the invention may be used to assist in refolding recombinantly produced polypeptides, which are obtained in an unfolded or misfolded form. Thus, recombinantly produced polypeptides may be passed down a column on which is immobilised a composition comprising protein disulphide isomerase and/or a molecular chaperone and/or a prolyl peptidyl isomerase.

In an alternative embodiment, in a the invention may be employed to maintain the folded conformation of proteins, for example during storage, in order to increase shelf life. under storage conditions, many proteins lose their activity, as a result of disruption of correct folding. The presence of molecular chaperones, in combination with foldases, reduces or reverses the tendency of polypeptides to become unfolded and thus greatly increases the shelf life thereof. In this embodiment, the invention may be applied to reagents which comprise polypeptide components, such as enzymes, tissue culture components, and other proteinaceous reagents stored in solution.

In a further embodiment, the invention may be used to potentiate the correct folding of proteins which, through storage, exposure to denaturing conditions or otherwise, have become misfolded. Thus, the invention may be used to recondition reagents or other proteins. For example, proteins in need of reconditioning may be passed down a column to which is immobilised a combination of reagents in accordance with he invention.

Alternatively, beads having immobilised thereon such a combination may be suspended in a solution comprising the proteins in need of reconditioning. Moreover, the components of the combination according to the invention may be added in solution to the proteins in need of reconditioning.

As noted above, the components of the combination according to the invention may comprise derivatives of molecular chaperones or foldases, including variants of such polypeptides which retain common structural features thereof. Variants which retain common structural features can be fragments of molecular chaperones or foldases. Fragments of molecular chaperones or foldases comprise smaller polypeptides derived from therefrom. Preferably, smaller polypeptides derived from the molecular chaperones or foldases according to the invention define a single feature which is characteristic of the molecular chaperones or foldases. Fragments may in theory be almost any size, as long as they retain the activity of the molecular chaperones or foldases described herein.

With respect to molecular chaperones of the GroEL/hsp-60 family, a preferred set of fragments have been identified which possess the desired activity. These fragments are set forth in our copending international patent application PCT/GB96/02980 and in essence comprise any fragment comprising at least amino acid residues 230-271 of intact

10007344.110801

GroEL, or their equivalent in another hsp-60 chaperone. Preferably, the fragments should not extend beyond residues 150-455 or 151-456 of GroEL or their equivalent in another hsp-60 chaperone. Where the fragments are GroEL fragments, they must not possess the mutant GroEL sequence as set forth above; in other words, they must not have an Alanine residue at position 262 and/or an Isoleucine residue at position 267 of the sequence of intact GroEL.

Advantageously, the fragments comprise the apical domain of GroEL, or its equivalent in other molecular chaperones, or a region homologous thereto as defined herein. The apical domain spans amino acids 191-376 of intact GroEL. This domain is found to be homologous amongst a wide number of species and chaperone types.

Preferably, molecular chaperones according to the invention are homologous to, or are capable of hybridising under stringent conditions with, a region corresponding to the apical domain of GroEL as defined above.

In a highly preferred embodiment, the fragments are selected from the group consisting of residues 191-376, 191-345 and 191-335 of the sequence of intact GroEL.

Derivatives of the molecular chaperones or foldases also comprise mutants thereof, including mutants of fragments and other derivatives, which may contain amino acid deletions, additions or substitutions, subject to the requirement to maintain the activity of the molecular chaperones or foldases described herein. Thus, conservative amino acid substitutions may be made substantially without altering the nature of the molecular chaperones or foldases, as may truncations from the 5' or 3' ends. Deletions and substitutions may moreover be made to the fragments of the molecular chaperones or foldases comprised by the invention. Mutants may be produced from a DNA encoding a molecular chaperone or foldase which has been subjected to in vitro mutagenesis resulting e.g. in an addition, exchange and/or deletion of one or more amino acids. For example, substitutional, deletional or insertional variants of molecular chaperones or foldases can be prepared by recombinant methods and screened for immuno-crossreactivity with the native forms of the relevant molecular chaperone or foldase.

The fragments, mutants and other derivative of the molecular chaperones or foldases preferably retain substantial homology with the native molecular chaperones or foldases. As used herein, "homology" means that the two entities share sufficient characteristics for the skilled person to determine that they are similar in origin and function. Preferably, homology is used to refer to sequence identity. Thus, the derivatives of molecular chaperones or foldases preferably retain substantial sequence identity with native forms of the relevant molecular chaperone or foldase.

In the context of the present invention, a homologous sequence is taken to include an amino acid sequence which is at least 60, 70, 80 or 90% identical, preferably at least 95 or 98% identical at the amino acid level over at least 20, preferably 30 amino of the minichaperone. In particular, homology should typically be considered with respect to those regions of the sequence known to be essential for the chaperone activity rather than non-essential neighbouring sequences. Although homology can also be considered in terms of similarity (i.e. amino acid residues having similar chemical properties/functions), in the context of the present invention it is preferred to express homology in terms of sequence identity.

Homology comparisons can be conducted by eye, or more usually, with the aid of readily available sequence comparison programs. These commercially available computer programs can calculate % homology between two or more sequences.

% homology may be calculated over contiguous sequences, i.e. one sequence is aligned with the other sequence and each amino acid in one sequence directly compared with the corresponding amino acid in the other sequence, one residue at a time. This is called an "ungapped" alignment. Typically, such ungapped alignments are performed only over a relatively short number of residues (for example less than 50 contiguous amino acids).

Although this is a very simple and consistent method, it fails to take into consideration that, for example, in an otherwise identical pair of sequences, one insertion or deletion will cause the following amino acid residues to be put out of alignment, thus potentially resulting in a large reduction in % homology when a global alignment is performed. Consequently, most sequence comparison methods are designed to produce optimal alignments that take into

consideration possible insertions and deletions without penalising unduly the overall homology score. This is achieved by inserting "gaps" in the sequence alignment to try to maximise local homology.

- 5 However, these more complex methods assign "gap penalties" to each gap that occurs in the alignment so that, for the same number of identical amino acids, a sequence alignment with as few gaps as possible - reflecting higher relatedness between the two compared sequences - will achieve a higher score than one with many gaps. "Affine gap costs" are typically used that charge a relatively high cost for the existence of a gap and a smaller penalty for each subsequent residue in the gap. This is the most commonly used gap scoring system. High gap penalties will of course produce optimised alignments with fewer gaps. Most alignment programs allow the gap penalties to be modified. However, it is preferred to use the default values when using such software for sequence comparisons. For example when using the GCG Wisconsin Bestfit package (see below) the default gap penalty for amino acid sequences is -12 for a gap and -4 for each extension.

- Calculation of maximum % homology therefore firstly requires the production of an optimal alignment, taking into consideration gap penalties. A suitable computer program for carrying out such an alignment is the GCG Wisconsin Bestfit package (University of Wisconsin, U.S.A.; Devereux *et al.*, 1984, Nucleic Acids Research 12:387). Examples of other software than can perform sequence comparisons include, but are not limited to, the BLAST package (see <http://www.ncbi.nih.gov/BLAST/>), FASTA (Atschul *et al.*, 1990, J. Mol. Biol., 403-410; FASTA is available for online searching at, for example, <http://www2.ebi.ac.uk.fasta3>) and the GENWORKS suite of comparison tools.
- 25 However it is preferred to use the GCG Bestfit program.

- Although the final % homology can be measured in terms of identity, the alignment process itself is typically not based on an all-or-nothing pair comparison. Instead, a scaled similarity score matrix is generally used that assigns scores to each pairwise comparison based on chemical similarity or evolutionary distance. An example of such a matrix commonly used is the BLOSUM62 matrix - the default matrix for the BLAST suite of programs. GCG Wisconsin programs generally use either the public default values or a custom symbol comparison table if supplied (see user manual for further



details). It is preferred to use the public default values for the GCG package, or in the case of other software, the default matrix, such as BLOSUM62.

Once the software has produced an optimal alignment, it is possible to calculate %  
5 homology, preferably % sequence identity. The software typically does this as part of the sequence comparison and generates a numerical result.

Alternatively, sequence similarity may be defined according to the ability to hybridise to a complementary strand of a chaperone or foldase sequence as set forth above.

10 Preferably, the sequences are able to hybridise—with high stringency. Stringency of hybridisation refers to conditions under which polynucleic acids hybrids are stable. Such conditions are evident to those of ordinary skill in the field. As known to those of skill in the art, the stability of hybrids is reflected in the melting temperature ( $T_m$ ) of the hybrid  
15 which decreases approximately 1 to 1.5°C with every 1% decrease in sequence homology. In general, the stability of a hybrid is a function of sodium ion concentration and temperature. Typically, the hybridisation reaction is performed under conditions of higher stringency, followed by washes of varying stringency.

20 As used herein, high stringency refers to conditions that permit hybridisation of only those nucleic acid sequences that form stable hybrids in 1 M Na<sup>+</sup> at 65-68 °C. High stringency conditions can be provided, for example, by hybridisation in an aqueous solution containing 6x SSC, 5x Denhardt's, 1 % SDS (sodium dodecyl sulphate), 0.1 Na<sup>+</sup> pyrophosphate and 0.1 mg/ml denatured salmon sperm DNA as non specific competitor.

25 Following hybridisation, high stringency washing may be done in several steps, with a final wash (about 30 min) at the hybridisation temperature in 0.2 - 0.1x SSC, 0.1 % SDS.

Moderate stringency refers to conditions equivalent to hybridisation in the above described solution but at about 60-62°C. In that case the final wash is performed at the  
30 hybridisation temperature in 1x SSC, 0.1 % SDS.

Low stringency refers to conditions equivalent to hybridisation in the above described solution at about 50-52°C. In that case, the final wash is performed at the hybridisation temperature in 2x SSC, 0.1 % SDS.

- 5 It is understood that these conditions may be adapted and duplicated using a variety of buffers, e.g. formamide-based buffers, and temperatures. Denhardt's solution and SSC are well known to those of skill in the art as are other suitable hybridisation buffers (see, e.g. Sambrook, et al., eds. (1989) *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, New York or Ausubel, et al., eds. (1990) *Current Protocols in*  
10 *Molecular Biology*, John Wiley & Sons, Inc.). Optimal hybridisation conditions have to be determined empirically, as the length and the GC content of the probe also play a role.

- The invention also envisages the administration of polypeptide oligomers according to the invention as compositions, preferably for the treatment of diseases associated with protein  
15 misfolding. The active compound may be administered in a convenient manner such as by the oral, intravenous (where water soluble), intramuscular, subcutaneous, intranasal, intradermal or suppository routes or implanting (e.g. using slow release molecules). Depending on the route of administration, the active ingredient may be required to be coated in a material to protect said ingredients from the action of enzymes, acids and  
20 other natural conditions which may inactivate said ingredient.

- In order to administer the combination by other than parenteral administration, it will be coated by, or administered with, a material to prevent its inactivation. For example, the combination may be administered in an adjuvant, co-administered with enzyme inhibitors  
25 or in liposomes. Adjuvant is used in its broadest sense and includes any immune stimulating compound such as interferon. Adjuvants contemplated herein include resorcinols, non-ionic surfactants such as polyoxyethylene oleyl ether and n-hexadecyl polyethylene ether. Enzyme inhibitors include pancreatic trypsin.
- 30 Liposomes include water-in-oil-in-water CGF emulsions as well as conventional liposomes.

The active compound may also be administered parenterally or intraperitoneally. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

5 The pharmaceutical forms suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. In all cases the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable  
10 under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils. The proper fluidity can be maintained, for  
15 example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants.

The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic  
20 acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminium monostearate and gelatin.

25 Sterile injectable solutions are prepared by incorporating the active compound in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filtered sterilisation. Generally, dispersions are prepared by incorporating the sterilised active ingredient into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those  
30 enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and the freeze-drying technique which yield a powder of the active ingredient plus any additional desired ingredient from previously sterile-filtered solution thereof.

When the combination of polypeptides is suitably protected as described above, it may be orally administered, for example, with an inert diluent or with an assimilable edible carrier, or it may be enclosed in hard or soft shell gelatin capsules, or it may be compressed into tablets, or it may be incorporated directly with the food of the diet. For oral therapeutic administration, the active compound may be incorporated with excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. The amount of active compound in such therapeutically useful compositions in such that a suitable dosage will be obtained.

The tablets, troches, pills, capsules and the like may also contain the following: a binder such as gum tragacanth, acacia, corn starch or gelatin; excipients such as dicalcium phosphate; a disintegrating agent such as corn starch, potato starch, alginic acid and the like; a lubricant such as magnesium stearate; and a sweetening agent such as sucrose, lactose or saccharin may be added or a flavouring agent such as peppermint, oil of wintergreen, or cherry flavouring. When the dosage unit form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier.

Various other materials may be present as coatings or to otherwise modify the physical form of the dosage unit. For instance, tablets, pills, or capsules may be coated with shellac, sugar or both. A syrup or elixir may contain the active compound, sucrose as a sweetening agent, methyl and propylparabens as preservatives, a dye and flavouring such as cherry or orange flavour. Of course, any material used in preparing any dosage unit form should be pharmaceutically pure and substantially non-toxic in the amounts employed. In addition, the active compound may be incorporated into sustained-release preparations and formulations.

As used herein "pharmaceutically acceptable carrier and/or diluent" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents and the like. The use of such media and agents for pharmaceutical active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredient, use thereof in the

therapeutic compositions is contemplated. Supplementary active ingredients can also be incorporated into the compositions.

It is especially advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the mammalian subjects to be treated; each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the novel dosage unit forms of the invention are dictated by and directly dependent on (a) the unique characteristics of the active material and the particular therapeutic effect to be achieved, and (b) the limitations inherent in the art of compounding such as active material for the treatment of disease in living subjects having a diseased condition in which bodily health is impaired.

The principal active ingredients are compounded for convenient and effective administration in effective amounts with a suitable pharmaceutically acceptable carrier in dosage unit form. In the case of compositions containing supplementary active ingredients, the dosages are determined by reference to the usual dose and manner of administration of the said ingredients.

In a further aspect there is provided the combination of the invention as hereinbefore defined for use in the treatment of disease. Consequently there is provided the use of a combination of the invention for the manufacture of a medicament for the treatment of disease associated with aberrant protein/polypeptide structure. The aberrant nature of the protein/polypeptide may be due to misfolding or unfolding which in turn may be due to an anomalous e.g. mutated amino acid sequence. The protein/polypeptide may be destabilised or deposited as plaques e.g. as in Alzheimer's disease. The disease might be caused by a prion. A polypeptide-based medicament of the invention would act to renature or resolubilise aberrant, defective or deposited proteins.

The invention is further described below, for the purposes of illustration only, in the following examples.

## Examples

## 1. General Experimental Procedures

- 5 **Bacterial and bacteriophage strains.** The *E. coli* strains used in this study were: C41(DE3), a mutant of BL21(DE3) capable of expressing toxic genes (Miroux, B. & Walker, J. E. (1996) J. Mol. Biol. 260, 289-298); SV2 (B178*groEL4+*), SV3 (B178*groEL59*) and SV6 (B178*groEL673*): isogenic strains carrying temperature-sensitive alleles of *groEL*; SV1(=B178) (Georgopoulos, C., Hendrix, R. W., Casjens, S. R. & Kaiser, A. D. (1973) J. Mol. Biol. 76, 45-60), AI90 (*AgroEL::kan<sup>R</sup>*) [pBAD-EL] (Ivic, A., Olden, D., Wallington, E. J. & Lund, P. A. (1997) Gene 194, 1-8), and TG1 (Gibson, T. J. (1984) Ph.D. thesis, University of Cambridge, U.K). Bacteriophage  $\lambda$  b2cI (Zeilstra-Ryalls, J., Fayet, O., Baird, L. & Georgopoulos, C. (1993) J. Bacteriol. 175, 1134-1143) was used according to standard methods (Arber, W., Enquist, L., Hohn, B., Murray, N. E. & Murray, K. (1983) in Lambda II., ed. R. W. Hendrix, J. W. r., F. W. Stahl and R. A. Weisberg (Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.), pp. 433-466); plaque formation was assayed at 30 °C. T4D0, a derivative of bacteriophage T4 (Zeilstra-Ryalls, J., Fayet, O., Baird, L. & Georgopoulos, C. (1993) J. Bacteriol. 175, 1134-1143), was used according to standard methods (Karam, J. D. (1994) Molecular biology of bacteriophage T4. (American Society for Microbiology, Washington, DC)); plaque formation was assayed at 37 °C.

- Plasmid constructions.** Standard molecular biology procedures were used (Sambrook, J., Fritsch, E. F. & Maniatis, T. (1989) Molecular Cloning. A Laboratory Manual (Cold Spring Harbor Laboratory Press, N.Y.)). The schematic organisation of the plasmids used in this study is represented Figure 2. *Gp31* gene was PCR (polymerase chain reaction) amplified using two oligonucleotides 5' – C TTC AGA CAT ATG TCT GAA GTA CAA CAG CTA CC – 3' and 5' – TAA CGG CCG TTA CTT ATA AAG ACA CGG AAT AGC – 3' producing a 358 bp DNA using pSV25 (van der Vies, S., Gatenby, A. & Georgopoulos, C. (1994) Nature 368, 654-656) as template. The DNA sequence of a part of the mobile loop of Gp31 (residues 25 to 43) was removed by PCR, as described (Hemsley, A., Arnheim, N., Toney, M. D., Cortopassi, G. & Galas, D. J. (1989) Nucleic

Acids Res. 17, 6545-6551), using oligonucleotides 5' – **GGA GAA GTT CCT GAA CTG** – 3' and 5' – **GGA TCC GGC TTG TGC AGG TTC** – 3', creating a unique *Bam*H I site (bold characters). *GroEL* gene minichaperone (corresponding to the apical domain of GroEL, residues 191 to 376; (Zahn, R., Buckle, A. M., Perret, S., Johnson, C. M. J., Corrales, F. J., Golbik, R. & Fersht, A. R. (1996) Proc. Natl. Acad. Sci. U.S.A. 93, 15024-15029)) was amplified by PCR using oligonucleotides, containing a *Bam*H I site (underlined), 5' – TTC GGA TCC GAA GGT ATG CAG TTC GAC C – 3' and 5' – GTT GGA TCC AAC GCC GCC TGC CAG TTT C – 3' and cloned into the unique *Bam*H I site of pRSETA-Gp31Δloop vector, inserting minichaperone GroEL(191-376) in frame

into Gp31Δloop sequence. The single ring GroEL<sub>SR1</sub> mutant contains four amino acid substitutions (R452E, E461A, S463A. and V464A) into the equatorial interface of GroEL, which prevent the formation of double rings (Weissman, J. S., Hohl, C. M., Kovalenko, O., Kashi, Y., Chen, S., Braig, K., Saibil, H. R., Fenton, W. A. & Horwich, A. L. (1995) Cell 83, 577-587). The corresponding mutations were introduced into *groEL* by PCR (Hemsley, A., Arnheim, N., Toney, M. D., Cortopassi, G. & Galas, D. J. (1989) Nucleic Acids Res. 17, 6545-6551) using oligonucleotides 5' – TGA GTA CGA TCT GTT CCA GCG GAG CTT CC – 3' and 5' – ATT GCG GCG AAG CGC CGG CTG CTG TTG CTA ACA CCG – 3' and pRSETA-*Eag* I GroEL or GroESL vectors (Chatellier, J., Hill, F., Lund, P. & Fersht, A. R. (1998) Proc. Natl. Acad. Sci. U.S.A. 95, 9861-9866) as template; silent mutations, in respect to the codon usage in *E. coli*, create a unique *Mfe* I (bold characters) and *Nae* I (underlined). *GroEL*(E191G; *groEL44* allele) gene was PCR amplified from *E. coli* SV2 strain (Zeilstra-Ryalls, J., Fayet, O., Baird, L. & Georgopoulos, C. (1993) J. Bacteriol. 175, 1134-1143) using two oligonucleotides 5' – T AGC TGC CAT ATG GCA GCT AAA GAC GTA AAA TTC GG – 3' and 5' – ATG TAA CGG CCG TTA CAT CAT GCC GCC CAT GCC ACC – 3' producing a 1,659 bp DNA with unique sites for *Nde* I and *Eag* I (underlined). The different genes were subcloned into the unique *Nde* I and *Eag* I unique sites of pACYC184, pJC and pBAD30 (Guzman, L.-M., Belin, D., Carson, M. J. & Beckwith, J. (1995) J. Bacteriol. 177, 4121-4130) vectors (Chatellier, J., Hill, F., Lund, P. & Fersht, A. R. (1998) Proc. Natl. Acad. Sci. U.S.A. 95, 9861-9866). A colony-based PCR procedure was used to identify the positive clones (Chatellier, J., Mazza, A., Brousseau, R. & Vernet, T. (1995) Analyt. Biochem. 229, 282-290). PCR cycle sequencing using fluorescent dideoxy chain terminators (Applied Biosystems) were performed and analysed on an Applied

Biosystems 373A Automated DNA. All PCR amplified DNA fragments were sequenced after cloning.

**Proteins expression purification and characterisation.** The GroE proteins, ~57.5 kDa

5 GroEL and ~10 kDa GroES, were expressed and purified as previously described (Chatellier, J., Hill, F., Lund, P. & Fersht, A. R. (1998) Proc. Natl. Acad. Sci. U.S.A. 95, 9861-9866; Corrales, F. J. & Fersht, A. R. (1996) Folding & Design 1, 265-273). GroEL<sub>SR1</sub> mutant was expressed and purified using the same procedure used for the wild-type GroEL; GroEL<sub>SR1</sub> mutant was separated from endogenous wild-type GroEL by  
10 ammonium sulphate precipitation at 30% saturation. GroEL(E191G) protein was expressed by inducing the P<sub>BAD</sub> promoter of pBAD30 based vector with 0.2 % arabinose in *E. coli* SV2 strain (Zeilstra-Ryalls, J., Fayet, O., Baird, L. & Georgopoulos, C. (1993) J. Bacteriol. 175, 1134-1143). Purification was performed essentially as described (Corrales, F. J. & Fersht, A. R. (1996) Folding & Design 1, 265-273). Residual peptides  
15 bound to GroEL proteins were removed by ion-exchange chromatography on a MonoQ column (Pharmacia Biotech.) in presence of 25 % methanol. The over-expression of histidine-tagged (short histidine tail; sht)-minichaperone GroEL(191-376) in *E. coli* C41(DE3) cells and, the purification and the removal of sht by thrombin cleavage were carried out essentially as previously described (Zahn, R., Buckle, A. M., Perret, S.,  
20 Johnson, C. M. J., Corrales, F. J., Golbik, R. & Fersht, A. R. (1996) Proc. Natl. Acad. Sci. U.S.A. 93, 15024-15029). Gp31 proteins wild-type (~12 kDa), Δloop (~10.4 kDa) and MC<sub>7</sub> (~30.6 kDa), were expressed by inducing the T7 promoter of pRSETA-Eag I based vectors with isopropyl-β-D-thiogalactoside (IPTG) in *E. coli* C41(DE3) (Miroux, B. & Walker, J. E. (1996) J. Mol. Biol. 260, 289-298) overnight at 25 °C. Purification  
25 procedures were essentially as described (van der Vies, S., Gatenby, A. & Georgopoulos, C. (1994) Nature 368, 654-656; Castillo, C. J. & Black, L. W. (1978) J. Biol. Chem. 253, 2132-2139). Ammonium sulphate precipitation (only 20% saturation for Δloop; 30 to 70% saturation for wild-type and MC<sub>7</sub>) was followed by ion-exchange chromatography on a DEAE-Sepharose column (Pharmacia Biotech.). Gp31 proteins were eluted with a 0-  
30 0.5 M NaCl gradient in 20 mM Tris-HCl, 1 mM EDTA, 1 mM β-mercaptoethanol, pH 7.5; Δloop and MC<sub>7</sub> eluted between 0.32-0.44 and 0.38-0.48 mM NaCl, respectively. Gp31 proteins were further purified by gel filtration chromatography on a Superdex™ 200 (Hiload 26/10) column (Pharmacia Biotech.) equilibrated with 100 mM Tris-HCl, pH



7.5 and, dialysed against and stored in 50 mM Tris-HCl, 0.1 mM EDTA, 1 mM  $\beta$ -mercaptoethanol, pH 7.5. Proteins were analysed by electrospray mass spectrometry. Protein concentration was determined by absorbance at 276 nm using the method of Gill & von Hippel (Gill, S. C. & von Hippel, P. H. (1989) *Analyt. Biochem.* 182, 319-326) and confirmed by quantitative amino acid analysis.

Constitutive expression under the control of the tetracycline-resistance gene promoter / operator was obtained using the high copy-number pJC vectors (Chatellier, J., Hill, F., Lund, P. & Fersht, A. R. (1998) *Proc. Natl. Acad. Sci. U.S.A.* 95, 9861-9866). pBAD30 vector allows inducible expression with 0.2-0.5 % arabinose controlled by the *P<sub>BAD</sub>* promoter and its regulatory gene, *araC* (Guzman, L.-M., Belin, D., Carson, M. J. & Beckwith, J. (1995) *J. Bacteriol.* 177, 4121-4130). The level of expression of MC<sub>7</sub> was analysed by 15% sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE) under non-reducing conditions followed by Western blotting as described (Chatellier, J., Hill, F., Lund, P. & Fersht, A. R. (1998) *Proc. Natl. Acad. Sci. U.S.A.* 95, 9861-9866).

**Molecular weight determination by analytical gel filtration chromatography and analytical ultracentrifugation.** One hundred  $\mu$ l aliquots of protein (1 mg.mL<sup>-1</sup>) were loaded onto a Superdex™ 200 (HR 10/30) column (Pharmacia Biotech.) equilibrated with 50 mM Tris-HCl, 150 mM NaCl, pH 7.5 at 0.5 mL.min<sup>-1</sup> at 20 °C. The column was calibrated using gel filtration standards from Pharmacia Biotech. (thyroglobulin, MW=669 kDa; ferritin, MW=440 kDa; aldolase, MW=158 kDa; ovalbumin, MW=45 kDa; chymotrypsinogen MW=25 kDa; RNase, MW=13 kDa). Molecular weights were determined by logarithmic interpolation.

Sedimentation analysis was performed in 50 mM Tris-HCl, 2.5 mM DTE (dithioerythritol), pH 7.2 at 20 °C with protein concentration in the range 45-300  $\mu$ M, scanning at 280 nm, with a Beckman XL-A analytical ultracentrifuge, using an An-60Ti rotor. Sedimentation equilibrium experiments were at 10,000 rev.min<sup>-1</sup> with overspeeding at 15,000 rev.min<sup>-1</sup> for 6 hours to speed the attainment of equilibrium. Scans were taken at intervals of 24 hours, until successive scans superimposed exactly, when the later scan

was taken as being operationally at equilibrium. To evaluate the apparent average molecular weight, data were fitted by non-linear regression.

**Circular dichroism spectroscopy (CD).** Far UV (200-250 nm)-CD spectra at 25 °C were measured on a Jasco J720 spectropolarimeter interfaced with a Neslab PTC-348WI water bath, using a thermostatted cuvette of 0.1 cm path length. Spectra are averages of 10 scans and were recorded with a sampling interval of 0.1 nm. Thermal denaturation was carried out from 5-95 °C at a linear rate of 1 °C.min<sup>-1</sup> and monitored at 222 nm. The reversibility was checked after incubation at 95 °C for 20 min and cooling to and equilibration at 5 °C. The protein concentration was 45 µM in 10 mM sodium phosphate buffer pH 7.8, 2.5 mM DTE (dithioerythrol).

**GroES binding and competition assays by ELISA** (enzyme-linked immunosorbant assay). Proteins were coated onto plastic microtitre plates (Maxisorb, Nunc) overnight at 4 °C at a concentration of 10 µg/mL in carbonate buffer (50 mM NaHCO<sub>3</sub>, pH 9.6). Plates were blocked for 1 hour at 25 °C with 2% Marvel in PBS (phosphate buffered saline: 25 mM NaH<sub>2</sub>PO<sub>4</sub>, 125 mM NaCl, pH 7.0). GroES, at 10 µg/mL in 100 µL of 10 mM Tris-HCl, 200 mM KCl, pH 7.4, were bound at 25 °C for 1 hour. Bound GroES were detected with rabbit anti-GroES antibodies (Sigma) followed by anti-rabbit immunoglobulins horseradish peroxidase conjugated antibodies (Sigma).

A peptide corresponding to the mobile loop of GroES (residues 16 to 32, numbered as in Hemmingsen, S. M., Woolford, C., van, d. V. S., Tilly, K., Dennis, D. T., Georgopoulos, C. P., Hendrix, R. W. & Ellis, R. J. (1988) *Nature* 333, 330-334) was synthesised as described (Chatellier, J., Buckle, A. M. & Fersht, A. R. (1999) *J. Mol. Biol.*, in press). The inhibition of the binding of MC<sub>7</sub> proteins by the free peptide was analysed by ELISA, essentially as above, by adding different concentrations (between 10,000 to 0.1 µM) of free peptide solved in 0.1% TFA solution to 1 µg of proteins prior incubation to coated GroES proteins (10 µg/mL). GroEL molecules were detected with rabbit anti-GroEL antibodies (Sigma) followed by anti-rabbit immunoglobulins horseradish peroxidase conjugate antibodies (Sigma). ELISAs were developed with 3',3',5',5'-tetramethylbenzidine (TMB, Boehringer Mannheim). Reactions were stopped with 50 µl

of 1M H<sub>2</sub>SO<sub>4</sub> after 10 min and readings taken by subtracting the O.D.<sub>650 nm</sub> from the O.D.<sub>450 nm</sub>.

**Anti-GroEL antibodies binding by ELISA.** The same amount of proteins (1 µg) were coated as described above. GroEL molecules were detected with either (i) rabbit anti-GroEL horseradish peroxidase conjugate antibodies (9 mg/mL; Sigma) or (ii) rabbit anti-GroEL antibodies (11.5 mg/mL; Sigma) followed by anti-rabbit immunoglobulins horseradish peroxidase conjugate antibodies (Sigma). ELISAs were developed as described above.

**In vitro refolding experiments.** Refolding assays of pig heart mitochondrial malate dehydrogenase (mtMDH; Boehringer-Mannheim) and aggregation protection were carried out essentially as described (Peres Ben-Zvi, A. P., Chatellier, J., Fersht, A. R. & Goloubinoff, P. (1998) Proc. Natl. Acad. Sci. U.S.A. 95, 15275-15280). The concentrations of MC<sub>7</sub> used were between 8-16 µM (reporter to protomer).

**In vivo complementation experiments.** Complementation experiments were performed by transforming electro-competent SV2 or SV6 cells with the pJC series of expression vectors and plating an aliquot of the transformation reactions directly at 43 °C. The percentage of viable cells relative to the growth at 30 °C was determined. A representative number of clones which grew at 43 °C were incubated in absence of any selective markers at permissive temperature. After prolonged growth the loss of the pJC plasmids and the ts phenotype were verified. Each experiment was performed in duplicate. Plasmids carrying no *groE* genes or encoding the GroE proteins were used as negative or positive controls, respectively.

P1 transduction (Miller, J. H. (1972) Experiments in Molecular Genetics (Cold Spring Harbor, N.Y.)), using strain AI90 (*ΔgroEL::kan<sup>R</sup>*) [pBAD-EL] as donor (Ivic, A., Olden, D., Wallington, E. J. & Lund, P. A. (1997) Gene 194, 1-8), was used to delete the *groEL* gene of TG1 cells transfected by the different pJC vectors. Transductants were selected on LB plates containing 10 µg/mL of kanamycin at 37 °C. Approximately 25 colonies were transferred onto plates containing kanamycin at 50 µg/mL. After incubation for 24 h at 37 °C, colonies that grew were screened by PCR as described.

AI90 (*AgroEL::kan<sup>R</sup>*) [pBAD-EL] cells were transformed with the pJC vector series. Transformants were selected at 37 °C on LB supplemented with 50 µg.mL<sup>-1</sup> of kanamycin, 120 µg.mL<sup>-1</sup> of ampicillin, 25 µg.mL<sup>-1</sup> of chloramphenicol and 0.2% L(+)-arabinose.

5 Depletion of GroEL protein was analysed at 37 °C by plating the same quantity of AI90 [pBAD-EL + pJC vectors] cells on LB plates containing 1% D(+)-glucose or various amount of arabinose.

Each experiment was performed in triplicate. Plasmids carrying no *groE* genes or encoding the GroE proteins were used as negative or positive controls, respectively.

**Effect on Lorist6 replication of over-expressing of MC<sub>7</sub>.** The effect of over-expressing Gp31 proteins from pJC vector series on the replication of the bacteriophage λ origin vector, Lorist6 (Gibson, T. J., Rosenthal, A. & Waterston, R. H. (1987) Gene 53, 283-286) in TG1 (Gibson, T. J. (1984) Ph.D. thesis, University of Cambridge, U.K) or SV1 (Georgopoulos, C., Hendrix, R. W., Casjens, S. R. & Kaiser, A. D. (1973) J. Mol. Biol. 76, 45-60) cells was determined essentially as described (Chatellier, J., Hill, F., Lund, P. & Fersht, A. R. (1998) Proc. Natl. Acad. Sci. U.S.A. 95, 9861-9866).

**2. Example 1: Gp31 protein as a scaffold for displaying heptameric GroEL minichaperone.** We describe a scaffold on which any polypeptide may be hung; as a result, the polypeptide is oligomerised. The scaffold is the bacteriophage T4 Gp31 (gene product) heptamer. The monomeric protein is 12 kDa, but it spontaneously forms a stable heptameric structure (90 kDa) of which the three-dimensional structure is known from X-ray crystallography (Hunt, J. F., van der Vies, S., Henry, L. & Deisenhofer, J. (1997) Cell 90, 361-371). This illustrates that a highly mobile polypeptide loop (residues 25 to 43; Chatellier, J., Mazza, A., Brousseau, R. & Vernet, T. (1995) Analyt. Biochem. 229, 282-290) projects from each subunit (Figure 1). The basis of the method is the substitution of this loop by a chosen peptide sequence.

In an effort to increase the avidity of minichaperones for substrates, and consequently to improve their chaperonin-facilitated protein folding, we generated the fusion protein, Gp31Δloop::GroEL(191-376) (hereafter named MC<sub>7</sub>), where the mobile loop of Gp31

was replaced by the sequence of minichaperone GroEL (residues 191 to 376) (Figure 2). MC<sub>7</sub> was cloned downstream of the T7 promoter of pRSETAsht-*Eag* I vector (Chatellier, J., Hill, F., Lund, P. & Fersht, A. R. (1998) Proc. Natl. Acad. Sci. U.S.A. 95, 9861-9866). After sonication, the soluble and insoluble fractions of IPTG-induced transfected C41(DE3) cells (Miroux, B. & Walker, J. E. (1996) J. Mol. Biol. 260, 289-298) were analysed by SDS-PAGE. Most of MC<sub>7</sub> was present in the insoluble fraction. Insoluble material dissolved in 8 M urea was efficiently refolded by dialysis at 4 °C. MC<sub>7</sub> was purified by ion-exchange and gel filtration chromatography. MC<sub>7</sub> was over-expressed in C41(DE3) cells to give 0.25-0.5 g purified protein per L of culture. Purified MC<sub>7</sub> coincided to seven 30.6 kDa subunits of Gp31Δloop::GroEL(191-376) as determined by analytical size exclusion chromatography (Figure 3 a) and analytical ultracentrifugation (Figure 3 b); Gp31Δloop corresponds to a tetra-decamer (14 subunits). The introduction of a foreign polypeptide in the Gp31 scaffold does prevent its oligomerisation ability. The electronic microscopy studies of MC<sub>7</sub> revealed views that correspond to front views of oligomers with a diameter close to the one of GroEL (J.L. Carrascosa, J.C. & A.R.F., unpublished). The circular dichroism spectrum of MC<sub>7</sub> indicated significant α-helical structure (Figure 4a). The thermal unfolding monitored by far UV-CD was reversible although more than one transition exist (Figure 4b).

Bacterial GroES or the human mitochondrial Hsp10 homologous oligomerisable scaffolds have been also successfully used to oligomerise polypeptides displayed in their mobile loops<sup>1</sup>.

### 3. Example 2: Binding to heptameric bacterial co-chaperonin, GroES.

The functionality of MC<sub>7</sub> was examined for binding to GroES, since the interaction between GroEL and GroES is known to be less favourable for one monomer than for the heptamer. MC<sub>7</sub> bound specifically to GroES, conversely monomeric minichaperone GroEL(191-376) did not detectably bind the bacterial co-chaperonin (Figure 5a).

---

<sup>1</sup> Please send me details of these experiments.

The ability of a synthetic peptide corresponding to residues 16 to 32 of GroES mobile loop to displace bound GroES from MC<sub>7</sub> was tested by competition ELISA. The synthetic GroES mobile loop peptide did inhibit the binding of MC<sub>7</sub> with an IC<sub>50</sub> of 10 µM compared to 100 µM for GroEL (Figure 5b). The apparent dissociation constant for the formation of the GroEL-GroES complex is low (10<sup>-6</sup> M), which is compatible with cycling of GroES on and off GroEL during chaperonin-assisted folding. On the other hand, GroEL<sub>SR1</sub> (Weissman, J. S., Rye, H. S., Fenton, W. A., Beechem, J. M. & Horwich, A. L. (1996) Cell 84, 481-490) is unable to release GroES in the absence of signal transmitted via the binding of ATP to an adjacent ring. The 10-fold decrease of the affinity of MC<sub>7</sub> for GroES may be sufficient for multiple binding and release cycles.

***In vitro* activity of MC<sub>7</sub>.** *In vitro*, heat- and dithiothreitol-denatured mitochondrial malate dehydrogenase (mtMDH) refolds in high yield only in the presence of GroEL, ATP, and the co-chaperonin GroES (Peres Ben-Zvi, A. P., Chatellier, J., Fersht, A. R. & Goloubinoff, P. (1998) Proc. Natl. Acad. Sci. U.S.A. 95, 15275-15280). Monomeric minichaperone GroEL(191-376) binds denatured mtMDH, protecting is aggregation (Figure 7a) but, it is ineffective in enhancing the refolding rate (Figure 7b). Conversely, MC<sub>7</sub>, which protect further denatured mtMDH from aggregation (Figure 7a) is active in refolding denatured mtMDH (Figure. 7a) with a rate of 0.02 nM.min<sup>-1</sup>, compared to 0.04 nM.min<sup>-1</sup> for wild-type GroEL alone (Figure 7b). After 120 min, the yield of refolded mtMDH by MC<sub>7</sub> is about 2.5-3 nM, compared to 6 nM of enzyme rescued by wild-type GroEL (Figure 7c). Although saturating concentration of GroES (4 µM) does increase about 3- to 5-fold the rates at the beginning of the refolding reaction, a 10-fold decrease of the final yield was observed; indicating the absence of multiple cycles of binding and release of GroES to MC<sub>7</sub> (data not shown). Nevertheless, MC<sub>7</sub> is more efficient than GroEL<sub>SR1</sub> mutant (Llorca, O., Pérez-Pérez, J., Carrascosa, J., Galan, A., Muga, A. & Valpuesta, J. (1997) J. Biol. Chem. 272, 32925-32932; Nielsen, K. L. & Cowan, N. J. (1998) Molecular Cell 2, 1-7; this study); remarkably, MC<sub>7</sub> is only 2-fold less active than wild-type GroEL in refolding a non-permissive substrate *in vitro*.

4. **Example 3: *In vivo* complementation of thermosensitive *groEL* mutant alleles at 43 °C.**

We sought complementation of two thermosensitive (*ts*) *groEL* mutants of *E. coli* at 43 °C. *E. coli* SV2 has the mutation Glu191→Gly in GroEL corresponding to *groEL44* allele, while SV6 carries the *EL673* allele, which has two mutations, Gly173→Asp and Gly337→Asp. Complementation experiments were performed by transforming the thermosensitive (*ts*) *E. coli* strains SV2 or SV6 with the pJC series of expression vectors (Chatellier, J., Hill, F., Lund, P. & Fersht, A. R. (1998) Proc. Natl. Acad. Sci. U.S.A. 95, 9861-9866) and plating an aliquot of the transformation reaction directly at 43 °C. Subsequently, plasmids pJC from a representative number of individual clones growing at 43 °C were lost in the absence of continued chloramphenicol selection. Nearly all (≥ 95%) the cured clones were thermosensitive at 43 °C indicating the absence of recombination events for the reconstitution of wild-type *groEL* gene. The results obtained are qualitatively similar to those previously described (Chatellier, J., Hill, F., Lund, P. & Fersht, A. R. (1998) Proc. Natl. Acad. Sci. U.S.A. 95, 9861-9866). Only minichaperone sht-GroEL(193-335) complements the defect in SV2. The defective *groEL* in SV6 was complemented by expression of minichaperone sht-GroEL(191-345), and less well by sht-GroEL(193-335). Conversely, MC<sub>7</sub> and GroEL<sub>SR1</sub> complement both temperature-sensitive *E. coli groEL44* and *groEL673* alleles at 43 °C (Table 1). Colony-forming units were not observed for either strain at 43 °C with vectors either lacking inserts (pJCsh) or lacking GroEL(191-376) (pJCGp31Δloop).

It has been suggested the higher stability of shortest minichaperone sht-GroEL(193-335) could be responsible for the complementation of *groEL44* mutant allele. To test this eventuality, we purified GroEL(E191G; *groEL44* allele) mutant and compared its thermal stability with the wild-type GroEL. We found no difference in stability between the mutant and the wild-type proteins in presence or absence of ATP. In addition, highly stable functional mutants of GroEL (193-345) do not complement, as the parental minichaperone (Table 1), the defects in SV2 or even SV6. We concluded the thermal stability of minichaperone is not accountable for the complementation of *groEL* defects.

**Table 1.** Relative colony forming ability of transformed *ts groEL44* or *groEL673* *E. coli* strains at 43 °C.

Plasmids pJC	GroEL strains	
	SV2 <i>groEL44</i>	SV6 <i>groEL673</i>
short his tag (sht) ( <i>ES<sup>-</sup></i> , <i>EL<sup>-</sup></i> )	< 10 <sup>-4</sup>	< 10 <sup>-4</sup>
GroES(1-97)	5 x 10 <sup>-3</sup>	< 10 <sup>-4</sup>
Gp31(1-111)	0.5 x 10 <sup>-3</sup>	< 10 <sup>-4</sup>
Gp31 Δloop	< 10 <sup>-4</sup>	< 10 <sup>-4</sup>
GroEL(1-548)	1	1
GroES-EL	1	1
sht-GroEL(191-376)	< 10 <sup>-4</sup>	< 10 <sup>-4</sup>
sht-GroEL(191-345)	0.01-0.02	0.07-0.09
sht-GroEL(193-335)	0.05-0.09	0.03-0.05
Gp31Δ::GroEL(191-376)	0.15-0.2	0.1

5

#### 5. Example 4: *In vivo* complementation at 37 °C.

The effects of MC<sub>7</sub> on the growth at 37 °C of a strain of *E. coli* in which the chromosomal *groEL* gene had been deleted were analysed in two ways. First, we attempted to delete the *groEL* gene of TG1 which had been transformed with the different pJC MC<sub>7</sub> vector by P1 transduction. However, no transductants could be obtained where the *groEL* gene had been deleted, unless intact GroEL was expressed from the complementing plasmid. This is consistent with the known essential role of GroEL. Second, we analysed the complementation of AI90 (*ΔgroEL::kan<sup>R</sup>*) [pBAD-EL] *E. coli* strain. In this strain, the chromosomal *groEL* gene has been deleted and GroEL is expressed exclusively from a plasmid-borne copy of the gene which can be tightly regulated by the arabinose *P<sub>BAD</sub>* promoter and its regulatory gene, *araC*. AraC protein acts as either a repressor or an activator depending on the carbon source used. *P<sub>BAD</sub>* is activated by arabinose but repressed by glucose (Guzman, L.-M., Belin, D., Carson, M. J. & Beckwith, J. (1995) J.



Bacteriol. 177, 4121-4130). The AI90 [pBAD-EL] cells can not grow on medium supplemented with glucose at 37 °C (Ivic, A., Olden, D., Wallington, E. J. & Lund, P. A. (1997) Gene 194, 1-8). As minichaperones (Chatellier, J., Hill, F., Lund, P. & Fersht, A. R. (1998) Proc. Natl. Acad. Sci. U.S.A. 95, 9861-9866.), MC<sub>7</sub> was unable to suppress this *groEL* growth defect (Table 2). We then determined whether MC<sub>7</sub> could supplement low levels of GroEL from transfected AI90 [pBAD-EL]. At 0.01% arabinose, cells transfected with pJC expressing *sht* alone, Gp31Δloop or *sht*-GroEL(191-376), showed little colony forming ability (less than 5%). But those containing pJC MC<sub>7</sub> produced about 30% of the number produced in the presence of 0.2% arabinose. Thus, pJC MC<sub>7</sub>, but not pJCGroEL<sub>SR1</sub>, significantly supplements depleted levels of GroEL, about twice as pJC *sht*-GroEL(193-335) (Chatellier, J., Hill, F., Lund, P. & Fersht, A. R. (1998) Proc. Natl. Acad. Sci. U.S.A. 95, 9861-9866).

**Table 2.** Plating ability of transformed AI90 ( $\Delta groEL::kan^R$ ) [pBADEL] *E. coli* strain at 37 °C in presence of different amount of arabinose.

Plasmids pJC	% L(+)arabinose			
	0.15	0.10	0.01	0.00
short his tag ( <i>sht</i> ) ( <i>ES</i> <sup>-</sup> , <i>EL</i> <sup>-</sup> )	++	+	+/-	-
Gp31Δloop	++	+	+/-	-
GroEL (1-548)	+++	+++	+++	+++
<i>sht</i> -GroEL (191-376)	++	+	+/-	-
Gp31Δ:: GroEL (191-376)	+++	+++	+	-
+++ , growth identical to that in presence of 0.2 % L(+)arabinose (100 %), in terms of both number and size; ++, about 50 % of the colonies relative to that in presence of 0.2				

% L(+)*arabinose*; +, about 30 % of the colonies; +/-,  $\leq 5$  % of the colonies and size reduced relative to that in presence of 0.2 % L(+)*arabinose*; -, no visible colonies.

5    **6. Example 5: Effect on bacteriophages  $\lambda$  and T4 growth of over-expressing MC<sub>7</sub>.**

Bacteriophages  $\lambda$  and T4 require the chaperonins GroES and GroEL for protein folding during morphogenesis (Zeilstra-Ryalls, J., Fayet, O. & Georgopoulos, C. (1991) Annu. Rev. Microbiol. 45, 301-325). Nine *groE* alleles which fail to support  $\lambda$  growth have been  
10 sequenced (Zeilstra-Ryalls, J., Fayet, O., Baird, L. & Georgopoulos, C. (1993) J. Bacteriol. 175, 1134-1143). We examined the ability of MC<sub>7</sub>, over-expressed from the constitutive tet promoter on a high-copy number vector (see Figure 2), to complement three mutant *groEL* alleles for plaque formation by  $\lambda$  (b2cI) at 30 °C (Table 3) and T4 at 37 °C (Table 4).

15    The *groE* operon was named for its effects on the E protein of  $\lambda$  (Georgopoulos, C., Hendrix, R. W., Casjens, S. R. & Kaiser, A. D. (1973) J. Mol. Biol. 76, 45-60). Although heat induction of the *groE* operon has been shown to decrease burst size of  $\lambda$  bacteriophage in *E. coli* (Wegrzyn, A., Wegrzyn, G. & Taylor, K. (1996) Virology 217, 594-597). In contrast, we showed that the over-expression of GroEL alone, which  
20 resulted in slower growth of the bacteria, suffices to inhibit  $\lambda$  growth (Table 3). This effect was specific; over-expression of GroEL together with GroES caused only a four-fold drop in plaques. Over-expression of GroES alone had no effect. Minichaperone GroEL(191-376) had no effect on plaque counts in SV1 (*groE*<sup>+</sup>). Conversely, over-  
25 expression of MC<sub>7</sub> prevents plaque formation by bacteriophage  $\lambda$  in SV1, but less markedly than GroEL (Table 3). It seems that the main effect of GroEL over-expression is mediated through the  $\lambda$  origin, which requires two proteins, O and P. As with GroEL, MC<sub>7</sub> (or GroEL<sub>SR1</sub>) inhibit the replication of the Lorist6 plasmid which use the bacteriophage  $\lambda$  origin. The effect on Lorist6 shows that the unfoldase activity is also an  
30 essential part of GroEL activity *in vivo*. MC<sub>7</sub> and minichaperones possess both, un- and folding, activities. GroEL over-expression gives weak complementation of  $\lambda$  growth in SV2 (*groEL44*) and SV3 (*groEL59*; Ser201→Phe). MC<sub>7</sub> does not, but GroEL<sub>SR1</sub> does

complement any of the *E. coli groEL* mutant strains for bacteriophage  $\lambda$  growth at 30 °C (Table 3).

Bacteriophage T4 (T4D0) also requires a functional *groEL* gene, but encodes a protein Gp31 which can substitute for GroES. The requirement for GroEL can be distinguished genetically from  $\lambda$ 's requirement. Thus only two of the four *groEL* alleles fail to support T4 replication; these are also the two thermosensitive mutations *EL44* and *EL673* (Zeilstra-Ryalls, J., Fayet, O., Baird, L. & Georgopoulos, C. (1993) *J. Bacteriol.* 175, 1134-1143; Zeilstra-Ryalls, J., Fayet, O. & Georgopoulos, C. (1991) *Annu. Rev. Microbiol.* 45, 301-325). While over-expression of Gp31 allows T4 growth in all strains (only SV2 and SV6 strains normally do not allow T4 growth), over-expression of Gp31 $\Delta$ loop inhibits T4 replication. On the other hand, MC<sub>7</sub> does, as does GroEL<sub>SR1</sub>, complement *E. coli groEL* mutant strains for bacteriophage T4 growth at 30 °C (Table 3).

**An unusual basis for thermo-sensitivity in a *groEL* mutation.** Surprisingly, over-expression of GroES demonstrates allele-specific complementation for  $\lambda$  and T4 of GroEL44 (Glu191→Gly) mutant (Tables 3 & 4). The effect is nevertheless incomplete; plaques on SV2 [pJCGroES] are invariably smaller than on SV1, or SV1 [pJCGroES]. The E191G single mutation blocks the assembly of the head structure of bacteriophage  $\lambda$ . A possible molecular basis for this allele-specificity lies in the nature of the *groEL44* mutation. The substitution of Glu191→Gly in the hinge region between the intermediate and apical domains of GroEL presumably increases the flexibility of the hinge, and thereby, modulates a hinged conformational change in GroEL required for proper interaction with GroES. Indeed, the pivoting of the hinge region ensures proper interaction with GroES. For example, the mutant GroEL59 (Ser201→Phe in the same hinge region) in SV3 has low affinity for GroES. Over-expression of GroES will favour the formation of GroES-EL44 complex; we indeed also observed complementation of SV2 for thermosensitivity and bacteriophages growth by over-expressing GroEL44 mutant. Taking advantage of the GroES effect, we observed that GroEL minichaperones and MC<sub>7</sub> all reduce both plaque size and number but, like GroEL, do not completely eliminate them in SV2 [pBADGroES].

GroEL44, purified to homogeneity, is effective in refolding heat- and DTT-denatured mitochondrial malate dehydrogenase in presence of ATP and saturating concentration of GroES. Surprisingly, GroEL44 is as thermo-stable as the wild-type GroEL, indicating the mutation does not destabilise the overall conformation of the mutant. As anticipated from our *in vivo* genetic analysis, the affinity between GroEL44 and GroES is decreased at 37 °C and even more at higher temperature.

Our results suggest that the *groEL44* mutation changes the distribution of GroEL subunits between apical domain-open and closed conformations. To allow GroEL<sub>SR1</sub> to release GroES in the absence of signal transmitted via the binding of ATP to an adjacent ring, we introduced the Glu191→Gly mutation in GroEL<sub>SR1</sub>, generating the GroEL<sub>SR2</sub> mutant. GroEL<sub>SR2</sub> is more efficient than MC<sub>7</sub> and even more than GroEL<sub>SR1</sub> *in vitro* and *in vivo*.

**Table 3.** Growth of bacteriophage  $\lambda$  at 30 °C in transformed wild-type and *groEL* mutant strains.

Plasmids pJC	<i>groEL</i> strains			
	SV1	SV2	SV3	SV6
	( <i>groEL</i> <sup>+</sup> )	<i>groEL44</i>	<i>GroEL59</i>	<i>groEL673</i>
short his tag (sht) ( <i>ES</i> <sup>-</sup> , <i>EL</i> <sup>-</sup> )	+++	-	-	-
GroES (1-97)	+++	+++	-	-
Gp31 (1-111)	+++	-	-	-
Gp31 $\Delta$ loop	+++	-	-	-
GroEL (1-548)	-	+	++	+/-
sht-GroEL (191-376)	+++	-	-	-
Gp31 $\Delta$ ::GroEL (191-376)	+	-	-	-

**Table 4.** Growth of bacteriophage T4 at 37 °C in transformed wild-type and *groEL* mutant strains.

Plasmids pJC	<i>groEL</i> strains			
	SV1 ( <i>groEL</i> <sup>+</sup> )	SV2 <i>groEL</i> 44	SV3 <i>groEL</i> 59	SV6 <i>groEL</i> 673
short his tag (sht) ( <i>ES</i> <sup>-</sup> , <i>EL</i> <sup>-</sup> )	+++	-	+++	-
GroES (1-97)	+++	+++	+++	-
Gp31 (1-111)	+++	+++	+++	+++
Gp31Δloop	+/-	-	-	-
GroEL (1-548)	+++	+++	+++	+++
sht-GroEL (191-376)	+++	-	+++	-
Gp31Δ::GroEL (191-376)	++	+	+	+/-

+++ , normal plaque-forming ability relative to wild-type *groEL*<sup>+</sup> strain, in terms of both number and size; ++, 5-fold fewer plaques relative to wild-type *groEL*<sup>+</sup> strain, or both; +, 10-fold fewer plaques, or plaque size reduced relative to wild-type *groEL*<sup>+</sup> strain, or both; +/-, 10<sup>2</sup>-fold fewer plaques and plaque size reduced relative to wild-type *groEL*<sup>+</sup> strain; -, no visible plaques (<10<sup>-4</sup>).

## 7. Example 6: MC<sub>72</sub>

A second oligomeric minichaperone polypeptide was constructed based on the GroES scaffold. This polypeptide, named MC<sub>72</sub>, is GroESΔloop::GroEL(191-376).

**Plasmid constructions** Standard molecular biology procedures were used (Sambrook et al., 1989). The plasmid pRSETA encoding GroES gene has been described (Chatellier et

al. 1998 In vivo activities of GroEL minichaperones. Proc. Natl. Acad. Sci. USA **95**, 9861-9866). The GroES mutant Gly24Trp was generated by polymerase chain reaction (PCR), as described (Hemsley et al., 1989 A simple method for site-directed mutagenesis using the polymerase chain reaction. Nucl. Acids Res. **17**, 6545-6551) using the template  
5 pRSETA encoding GroES (Chatellier et al., 1998) and the oligonucleotides 5' - C GGC TGG ATC GTT CTG ACC G - 3' and 5' - GC AGA TTT AGT TTC AAC TTC TTT ACG - 3', creating a Nae I site (bold characters).

The DNA sequence encoding a part of the mobile loop of GroES (residues 16 to 33) was  
10 removed by PCR, as described (Hemsley et al., 1989), using the oligonucleotides 5' - TCC GGC TCT GCA GCG G - 3' and 5' - TCC AGA GCC AGT TTC AAC TTC TTT ACG C - 3', creating a unique BamH I site (bold characters) and the vector pRSET A-GroESΔloop.

15 The GroEL minichaperone gene (corresponding to the apical domain of GroEL, residues 191 to 376; Zahn et al., 1996 Chaperone activity and structure of monomeric polypeptide binding domains of GroEL Proc. Nat. Acad. Sci. USA **93**, 15024-15029) was amplified by PCR and cloned into the unique BamH I site of pRSETA-GroESΔloop vector, thus inserting the minichaperone GroEL(191-376) in-frame into the GroESΔloop sequence.  
20 These genes were subcloned into the unique Nde I and Eag I sites of pACYC184, pJC and pBAD30 vectors (Guzman et al., 1995, Tight regulation, modulation, and high level expression by vectors containing the pBAD promoter. J. Bacteriol. **177**, 4121-4130; Chatellier et al., 1998). PCR cycle sequencing using fluorescent dideoxy chain terminators (Applied Biosystems) was performed and analysed on an Applied Biosystems  
25 373A machine. All PCR amplified DNA fragments were sequenced after cloning.

**Proteins expression, purification and characterization.** The GroES proteins, wild-type (~10.4 kDa) and mutant Gly24Trp (~10.5 kDa), Δloop (~9.8 kDa), MC<sub>72</sub> (~30 kDa), were  
30 expressed by inducing the T7 promoter of pRSETA-Eag I based vectors with isopropyl-β-D-thiogalactoside (IPTG) in E. coli C41(DE3) (Miroux & Walker, 1996 Over-production of proteins in Escherichia coli: Mutant hosts that allow synthesis of some membrane proteins and globular proteins at high levels. J. Mol. Biol. **260**, 289-298) overnight at 25 °C and purified as described (Chatellier et al., 1998).

Proteins were analysed by electrospray mass spectrometry. Protein concentration was determined by absorbance at 276 nm using the method of Gill & von Hippel (1989 Calculation of protein extinction coefficients from amino acid sequence data. Anal. Biochem. **182**, 319-326) and confirmed by quantitative amino acid analysis. In this study, protein concentrations refer to protomers, and not to oligomers.

**Characterisation of MC<sub>72</sub>** By both analytical size exclusion chromatography and analytical ultracentrifugation, both purified proteins, GroESΔloop and MC<sub>72</sub>, were heptamers of seven 9.8 and seven 30 kDa subunits, respectively. The introduction into the GroES scaffold of a foreign polypeptide substantially larger than itself did not prevent oligomerisation. Electron microscopic studies of MC<sub>72</sub> revealed a diameter close to that of GroEL.

**GroES binding** The functionality of MC<sub>72</sub> was verified by examining GroES binding (followed by fluorescence) and mtMDH refolding.

## 8:Example 7. Reduction of protein aggregation in Huntingdon's Disease

Huntington's disease (HD), spinocerebellar ataxias types 1 and 3 (SCA1, SCA3), and spinobulbar muscular atrophy (SBMA) are caused by CAG/polyglutamine expansion mutations (Perutz, M.F. 1999 Trend Biochem. Sci. **24**, 58-63; Rubinsztein, D.C. et al. 1999 J. Med. Genet. **36**, 265-270). A feature of these diseases is ubiquitinated intraneuronal inclusions derived from the mutant proteins, which colocalize with heat shock proteins (HSPs) in SCA1 and SBMA and proteasomal components in SCA1, SCA3, and SBMA. Previous studies suggested that HSPs might protect against inclusion formation, because overexpression of HDJ-2/HSDJ (a human HSP40 homologue) reduced ataxin-1 (SCA1) and androgen receptor (SBMA) aggregate formation in HeLa cells (See Wytenbach, A. et al. (2000) Proc. Natl. Acad. Sci. USA **97**, 2899-2903).

These phenomena have been studied by transiently transfecting part of Huntingdon exon 1 in COS-7, PC12, and SH-SY5Y cells. Inclusion formation was not seen with constructs



expressing 23 glutamines but was repeat length and time dependent for mutant constructs with 43–74 repeats. HSP70, HSP40, the 20S proteasome and ubiquitin colocalized with inclusions. Treatment with heat shock or with lactacystin, a proteasome inhibitor, increased the proportion of cells with inclusions of mutant huntingtin exon 1. Thus, inclusion formation may be enhanced in polyglutamine diseases, if the pathological process results in proteasome inhibition or a heat-shock response. Overexpression of HDJ-2/HSDJ did not modify inclusion formation in PC12 and SH-SY5Y cells but increased inclusion formation in COS-7 cells. To our knowledge, this is the first report of an HSP increasing aggregation of an abnormally folded protein in mammalian cells and expands the current understanding of the roles of HDJ-2yHSDJ in protein folding (See Wytenbach, A. et al. (2000) Proc. Natl. Acad. Sci. USA 97, 2899-2903).

In the eucaryotic cell, molecular chaperones might be involved in the actual formation of nuclear aggregates by stabilizing the unfolded protein in an intermediate conformation which has the propensity to interact with neighbouring, unfolded proteins (Chirmer, E.C. & Lindquist, S. 1997 Proc. Natl. Acad. Sci. USA 94: 13932-7; DebBurman, S.K. et al., 1997 Proc. Natl. Acad. Sci. . USA 94: 13938-43; Welch, W.J. & Gambetti, P. 1998 Nature 392: 23-4). The chaperone's dual roles in aggregate formation and suppression may not be mutually exclusive, but rather dependent on the presence and level of chaperone expression. For example, the yeast chaperone Hsp104 (or bacterial GroEL) was shown to be necessary, at intermediate levels, for the propagation of the prion-like factor [PSI<sup>+</sup>], but when Hsp104 was overexpressed, [PSI<sup>+</sup>] was lost. Overexpression of the yeast homologue Hsp70 also inhibited [PSI<sup>+</sup>] (Chernoff, Y.O. et al., 1995 Science 268: 880-4). A similar phenomenon may occur in spinocerebellar ataxia type 1, with endogenous levels of HDJ2/HDJ and/or Hsc70 contributing to the formation of ataxin-1 aggregates when the number of glutamine repeats is in the disease-causing range. As in yeast, it may be necessary to upregulate or modulate the level of molecular chaperones to reduce aggregate formation in affected neurons (Cummings, C.J. et al., 1998 Nat. Genet. 19: 148-54).

Recently, D. Rubinsztain et al. have shown that the overexpression of GroEL(191-345) minichaperone monomer reduced slightly but significantly the proportion of mutant

Huntingdon exon 1-expressing PC12 and SH-SY5Y cells with inclusions and also reduced cell death. We have tested MC<sub>72</sub> in the same system.

5 The overexpression of MC<sub>72</sub> [i.e. the fusion protein GroESΔloop::GroEL(191-376)], like yeast Hsp104, reduced inclusion formation and cell death even further in the same cells. On the other hand, the overexpression of wild-type GroEL alone, whose activity is regulated by its co-chaperone GroES and ATP hydrolysis, had no effect.

10 The failure of Hsps to release their substrates in polyQ disease may be a common feature indicating the use of chaperones as therapeutic agents in these cases.

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2

Then invention will now be further described by the following numbered paragraphs:

1. A polypeptide monomer capable of oligomerisation, said monomer comprising a polypeptide sequence which potentiates protein folding inserted into the sequence of a subunit of an oligomerisable protein scaffold.
2. A polypeptide monomer according to paragraph 1, wherein the oligomerisable protein scaffold subunit is selected from the group consisting of bacteriophage T4 Gp31, Escherichia coli GroES and homologues thereof of the cpn10 family.
3. A polypeptide monomer according to paragraph 1 or paragraph 2, wherein the polypeptide sequence is inserted into the sequence of the oligomerisable protein scaffold subunit such that both the N and C termini of the polypeptide monomer are formed by the sequence of the oligomerisable protein scaffold subunit.
4. A polypeptide monomer according to any preceding paragraph, wherein the polypeptide sequence is inserted into the oligomerisable protein scaffold subunit by replacing one or more amino acids thereof.
5. A polypeptide monomer according to paragraph 4, wherein the oligomerisable protein scaffold subunit is bacteriophage T4 Gp31 and the polypeptide sequence is inserted into the oligomerisable protein scaffold subunit by substantially replacing the mobile loop between amino acid positions 27 and 42.
6. A polypeptide monomer according to paragraph 4, wherein the oligomerisable protein scaffold subunit is Escherichia coli GroES and the polypeptide sequence is inserted into the oligomerisable protein scaffold subunit by substantially replacing the mobile loop between amino acid positions 19 and 29.
7. A polypeptide monomer according to paragraph 4, wherein the oligomerisable protein scaffold subunit is bacteriophage T4 Gp31 and the polypeptide sequence is inserted between positions 59 and 61 of the oligomerisable protein scaffold subunit.

8. A polypeptide monomer according to paragraph 4, wherein the oligomerisable protein scaffold subunit is Escherichia coli GroES and the polypeptide sequence is inserted between positions 56 and 57 of the oligomerisable protein scaffold subunit.
9. A polypeptide monomer according to paragraph 4, wherein the oligomerisable protein scaffold subunit is bacteriophage T4 Gp31 and polypeptide sequences are inserted at both positions described in paragraphs 5 and 7.
10. A polypeptide monomer according to paragraph 4, wherein the oligomerisable protein scaffold subunit is Escherichia coli GroES and polypeptide sequences are inserted at both positions described in paragraphs 6 and 8.
11. A polypeptide monomer according to paragraph 2, wherein the polypeptide sequence is displayed at the N or C terminus of the oligomerisable protein scaffold subunit.
12. A polypeptide oligomer comprising two or more polypeptide monomers according to any preceding paragraph.
13. A polypeptide oligomer according to paragraph 12, which is a homooligomer.
14. A polypeptide oligomer according to paragraph 12, which is a heterooligomer.
15. A polypeptide oligomer according to paragraph 14, wherein complementary protein folding are juxtaposed through the oligomerisation of different polypeptide monomers.
16. A polypeptide oligomer according to any one of paragraphs 12 to 15, wherein the monomers are covalently crosslinked.
17. A polypeptide oligomer according to any one of paragraphs 12 to 16, wherein the protein scaffold is in the form of a ring.
18. A polypeptide oligomer according to paragraph 17, wherein the ring is a heptameric ring.

19. A polypeptide monomer or oligomer according to any preceding paragraph, wherein the polypeptide sequence is selected from the group consisting of a minichaperone, a protease prosequence and a foldase.
20. A polypeptide oligomer or monomer according to paragraph 19, wherein the foldase is selected from the group consisting of a thiol/disulphide oxidoreductase and a peptidyl prolyl isomerase.
21. A method for promoting the folding of a polypeptide comprising contacting the polypeptide with a polypeptide oligomer or monomer according to paragraph 19 or paragraph 20.
22. A method according to paragraph 21, wherein the polypeptide is an unfolded or misfolded polypeptide.
23. A method according to paragraph 21 or 22, wherein the polypeptide comprises a disulphide.
24. A method according to any one of paragraphs 21 to 23, wherein the foldase is selected from the group consisting of thiol/disulphide oxidoreductases and peptidyl-prolyl isomerases.
25. A method according to paragraph 24, wherein the thiol/disulphide oxidoreductase is selected from the group consisting of E. coli DsbA and mammalian PDI, or a derivative thereof.
26. A method according to paragraph 24, wherein the peptidyl prolyl isomerase is selected from the group consisting of cyclophilin, parbulen, SurA and FK506 binding proteins.
27. A method according to any one of paragraphs 21 to 26, comprising contacting the polypeptide with a polypeptide oligomer according to any one of paragraphs 12 to 20 and a non-oligomerised foldase.

28. A method according to any one of paragraphs 21 to 27, wherein the polypeptide oligomer according to any one of paragraphs 21 to 26 and/or the foldase is immobilised onto a solid phase support.
29. A method according to paragraph 28 wherein the solid phase support is agarose.
30. A solid phase support having immobilised thereon a polypeptide oligomer according to any one of paragraphs 12 to 20 and/or a foldase.
31. A column packed at least in part with a solid phase support according to paragraph 30.
32. Use of a polypeptide according to any one of paragraphs 12 to 20, optionally in combination with a foldase, for promoting the folding of a polypeptide.
33. Use according to paragraph 32 wherein the polypeptide according to any one of paragraphs 12 to 20 and/or the foldase is immobilised on a solid phase support.
34. A composition comprising a combination of a polypeptide oligomer according to any one of paragraphs 12 to 20 and a foldase.